

4-3-2018

Diabetes and Obesity as Risk Factors for Surgical Site Infections in Laparoscopic Abdominal Surgery of Medicare Age Patients

FIU Herbert Wertheim College of Medicine Affiliates

Follow this and additional works at: <https://scholarlycommons.baptisthealth.net/se-all-publications>

Citation

FIU Herbert Wertheim College of Medicine Affiliates, "Diabetes and Obesity as Risk Factors for Surgical Site Infections in Laparoscopic Abdominal Surgery of Medicare Age Patients" (2018). *All Publications*. 2733.
<https://scholarlycommons.baptisthealth.net/se-all-publications/2733>

This Conference Poster -- Open Access is brought to you for free and open access by Scholarly Commons @ Baptist Health South Florida. It has been accepted for inclusion in All Publications by an authorized administrator of Scholarly Commons @ Baptist Health South Florida. For more information, please contact Carrie@baptisthealth.net.

Diabetes and Obesity as Risk Factors for Surgical Site Infections in Laparoscopic Abdominal Surgery of Medicare age patients

Andrew Manfra, MS; Arnav Gupta, MS; Israel Mostafa, MS; Grettel Castro; Juan Carlos Gonzales; Jaime Rodriguez; Noël C Barengo
 Medical and Population Health Sciences Research Department, Herbert Wertheim College of Medicine, Florida International University, Miami, FL, USA

Objective

The goal of this study was to determine if Diabetes Mellitus modifies the association between increasing Body Mass Index (BMI) and Surgical Site Infections in Medicare age patients who have undergone laparoscopic abdominal surgery.

Methods

Study Design: A retrospective cohort study with data extrapolated from the American College of Surgeons National Surgical Quality Improvement Program (ACS-NSQIP).

Study population: Study criteria were met by 54,064 patients >45y/o who underwent laparoscopic abdominal surgery in 2016. We excluded records of patients missing key variables or patients <45y/o to improve confounder distributions between Medicare age and older patients.

Independent Variables: Exposure was measured as the presence of DM, Medicare age >65y/o, and the presence of the classification of obesity (BMI>30) as per Center for Disease Control (CDC) BMI guidelines. BMI was calculated using patient height and weight on record and classified as BMI normal (<25 kg/m²), overweight (25-30 kg/m²), or obese (>30 kg/m²). Diabetic status was acquired by previous physician diagnosis as listed in patient charts and recorded as "Diabetes Mellitus requiring insulin or oral agents."

Outcome Variables: Any postoperative surgical site infection was recorded as an outcome event. Positive SSI in this study was defined if it met criteria defined by NSQIP as superficial incisional infection, deep incisional infection, or organ space infection.

Statistical analysis: Descriptive analysis was performed to check for missing variables. Bivariate analysis was performed according to the outcome and main exposure variable to check for possible confounders. Non-adjusted and adjusted logistic regression were used to assess the association between obesity, DM, and SSI.

Results

Table 1. Characteristics of study participants from NSQIP undergoing laparoscopic surgery of the abdomen according to body mass index category during 2016.

	BMI ^a Status				P-value
	Under (n=858) %	Normal (n=13,552) %	Overweight (n=18,979) %	Obese (n=18,979) %	
<u>Age(years)</u>					<0.001
45-65	76.87	82.47	88.4	94.1	
>65	23.13	17.5	11.6	5.9	
<u>Gender</u>					<0.001
Female	73.83	56.3	47.0	61.9	
Male	26.7	43.7	53.0	38.1	
<u>COPD^b</u>					<0.001
Yes	14.02	6.3	5.2	6.3	
no	85.98	93.7	94.9	93.7	
<u>History of CHF^c</u>					0.008
Yes	0.93	1.14	1.1	1.4	
No	99.07	98.86	98.9	98.6	
<u>Renal Failure post-op</u>					0.307
Yes	0.58	0.36	0.30	0.40	
no	99.42	99.64	99.70	99.6	
<u>Diabetes</u>					<0.001
Yes	5.84	11.3	16.8	28.4	
No	94.16	88.7	83.2	71.6	
<u>Previous sepsis</u>					<0.001
None	83.76	90.58	91.5	91.3	
SIRS ^d	0.64	4.89	4.8	5.0	
Septic	7.59	4.53	3.8	3.7	
<u>Emergency case</u>					<0.001
Yes	23.36	14.46	12.5	10.9	
No	76.64	85.54	87.5	89.1	

^aBody Mass Index; ^bChronic Obstructive Pulmonary Disease; ^cCongestive Heart Failure; ^dSystemic Inflammatory response syndrome. Emergency Cases defined as unplanned operations. Not listed (p<0.001); Ascites, Functional status, Hypertension, Disseminated Cancer. White Blood Cell Count. Smoking Status. Chronic Steroid use.

Table 2. Unadjusted and adjusted associations between BMI^a and Diabetes Mellitus

Characteristics	Surgical Site Infection			
	Unadjusted OR ^b	[95% Ci ^c]	Adjusted ^d OR	[95% CI]
<u>BMI</u>				
Normal	Ref.	Ref.	Ref.	Ref.
Underweight	1.63	[1.13-2.34]	1.29	[0.89-1.88]
Overweight	0.96	[0.83-1.11]	1.02	[0.88-1.18]
obese	1.23	[1.07-1.40]	1.29	[1.12-1.49]
<u>Diabetes</u>				
Yes	Ref.	Ref.	Ref.	
no	1.32	[1.17-1.50]	1.24	[1.08-1.41]

^aBody Mass Index; ^bOdds Ratio; ^cConfidence Interval; ^dAdjusted for age, sex, smoking, COPD hx, Ascites, Functional Status, CHF hx, Renal Failure, Hypertension, Disseminated Cancer, chronic steroid use, sepsis, emergency case, and White Blood Cell count.

Table 3. Adjusted associations between BMI^a and SSI^b according to Diabetes Mellitus Status

BMI	All		DM ^c (n=10,564)		No DM (n=43,108)	
	OR ^d	95%CI ^e	OR	95%CI	OR	95%CI
Normal	Ref	Ref	Ref	Ref	Ref	Ref
Under	1.30	[0.88-1.92]	1.11	[0.26-4.76]	1.30	[0.88-1.91]
Overweight	1.02	[0.87-1.20]	1.00	[0.69-1.45]	1.02	[0.87-1.20]
Obese	1.26	[1.08-1.48]	1.35	[0.96-1.90]	1.27	[1.08-1.49]

^aBody Mass Index; ^bSurgical Site Infection; ^cDiabetes Mellitus; ^dOdds Ratio; ^eConfidence Interval. Adjusted for age, sex, smoking, COPD hx, Ascites, Functional Status, CHF hx, Renal Failure, Hypertension, Disseminated Cancer, chronic steroid use, sepsis, emergency case, and WBC count.

Conclusion

- In this study we demonstrated that obesity and DM are independent risk factors for SSI in laparoscopic abdominal surgery.
- We found no evidence of effect modification by DM on obesity as a risk factor for SSI.
- The effect in our study when age was dichotomized to Medicare age was not statistically significant.
- We recommend surgeons not to use BMI as a predictor for SSI, but only when patients are obese BMI>30 kg/m² or have other comorbidities such as DM, CHF, COPD, disseminated cancer, and chronic steroid use.