

Baptist Health South Florida

## Scholarly Commons @ Baptist Health South Florida

---

All Publications

---

2018

### Associations between CT-determined visceral fat burden, hepatic steatosis, circulating white blood cell counts and neutrophil-to-lymphocyte ratio

Ricardo Cury

*Miami Cardiac & Vascular Institute, rcury@baptisthealth.net*

Follow this and additional works at: <https://scholarlycommons.baptisthealth.net/se-all-publications>

---

#### Citation

PLoS One (2018) 13(11):e0207284

This Article -- Open Access is brought to you for free and open access by Scholarly Commons @ Baptist Health South Florida. It has been accepted for inclusion in All Publications by an authorized administrator of Scholarly Commons @ Baptist Health South Florida. For more information, please contact [Carrief@baptisthealth.net](mailto:Carrief@baptisthealth.net).

RESEARCH ARTICLE


# Associations between CT-determined visceral fat burden, hepatic steatosis, circulating white blood cell counts and neutrophil-to-lymphocyte ratio

Kuo-Tzu Sung<sup>1,2,3</sup>, Richard Kuo<sup>1,2,4</sup>, Jing-Yi Sun<sup>5</sup>, Ta-Chuan Hung<sup>1,2,6</sup>, Shun-Chuan Chang<sup>1,2</sup>, Chuan-Chuan Liu<sup>6,7,8</sup>, Chun-Ho Yun<sup>1,2,4\*</sup>, Tung-Hsin Wu<sup>5</sup>, Chung-Lieh Hung<sup>1,2,3</sup>, Hung-I Yeh<sup>1,2,3</sup>, Charles Jia-Yin Hou<sup>1,2,3</sup>, Ricardo C. Cury<sup>9</sup>, David A. Zidar<sup>10</sup>, Hiram G. Bezerra<sup>10</sup>, Chris T. Longenecker<sup>10</sup>

**1** Department of Medicine, Mackay Medical College, Taipei, Taiwan, **2** Mackay Junior College of Medicine, Nursing and Management, Taipei, Taiwan, **3** Division of Cardiology, Department of Internal Medicine, Mackay Memorial Hospital, Taipei, Taiwan, **4** Department of Radiology, Mackay Memorial Hospital, Taipei, Taiwan, **5** Department of Biomedical Imaging and Radiological Sciences, National Yang Ming University, Taipei, Taiwan, **6** Department of Medical Technology, Yuanpei University of Science and Technology, Hsin-Chu, Taiwan, **7** Graduate Institute of Health Care Organization Administration, College of Public Health National Taiwan University, Taipei, Taiwan, **8** Health Evaluation Center, Mackay Memorial Hospital, Taipei, Taiwan, **9** Cardiovascular MRI and CT Program, Baptist Cardiac Vascular Institute, Miami, Florida, United States of America, **10** Division of Cardiology, Department of Internal Medicine, University Hospitals Harrington Heart & Vascular Institute, Case Western Reserve University, Cleveland, OH, United States of America

\* These authors contributed equally to this work.

\* [med202657@gmail.com](mailto:med202657@gmail.com)



## OPEN ACCESS

**Citation:** Sung K-T, Kuo R, Sun J-Y, Hung T-C, Chang S-C, Liu C-C, et al. (2018) Associations between CT-determined visceral fat burden, hepatic steatosis, circulating white blood cell counts and neutrophil-to-lymphocyte ratio. PLoS ONE 13(11): e0207284. <https://doi.org/10.1371/journal.pone.0207284>

**Editor:** Anna Halama, Weill Cornell Medical College in Qatar, QATAR

**Received:** June 19, 2018

**Accepted:** October 29, 2018

**Published:** November 20, 2018

**Copyright:** © 2018 Sung et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper and its Supporting Information files.

**Funding:** This work was supported by a research grant from Mackay Memorial Hospital and NSC 101-2314-B-195-020. There was no additional external funding received for this study.

**Competing interests:** The authors have declared that no competing interests exist.

## Abstract

Visceral adiposity is associated with cardiovascular disease, an association that may be mediated in part by inflammation. We hypothesized that regional measures of visceral adiposity would associate with commonly obtained clinical measures of immune status. We consecutively studied 3,291 subjects (mean age, 49.8±9.8 years) who underwent an annual cardiovascular risk survey. Peri-cardial (PCF) and thoracic peri-aortic adipose tissue (TAT) volumes were determined by dedicated computed tomography (CT) software (Aquarius 3D Workstation, TeraRecon, San Mateo, CA, USA). Hepatic steatosis was assessed by abdominal ultrasonography. We explored cross-sectional associations between visceral fat measures and high-sensitivity C-reactive protein (hs-CRP), leukocyte counts, and the neutrophil-to-lymphocyte ratio (NLR). Among 3,291 study participants, we observed positive linear associations between PCF and TAT, higher degree of hepatic steatosis and hs-CRP, various leukocyte counts, either total and its differential counts, and NLR (all trend  $p < 0.001$ ). Multi-variate linear and logistic regression models showed independent associations between PCF/TAT ( $\beta$ -Coef: 0.14/0.16, both  $p < 0.05$ ) and total WBC counts, with only TAT further demonstrated significant relations with neutrophil counts and NLR (both  $p < 0.05$ ) and independently identified abnormally high WBC and NLR (Odds ratio: 1.18 & 1.21, both  $p < 0.05$ ). C-statistics showed significant incremental model prediction for abnormally high WBC and NLR (both  $\Delta$ AUROC  $< 0.05$ ) when TAT was superimposed on traditional cardiovascular risks and biochemical information. Greater visceral adiposity burden and hepatic

**Abbreviations:** PCF, Pericardial adipose tissue; TAT, thoracic peri-aortic adipose tissue; VAT, visceral adipose tissue; MDCT, multi-detector computed tomography; hs-CRP, high-sensitive C-reactive protein; BMI, body mass index; BSA, body surface area; DBP, diastolic blood pressure; SBP, systolic blood pressure; eGFR, estimated glomerular filtration rate; HbA1c, glycosylated hemoglobin level; HDL, high-density lipoprotein; LDL, low-density lipoprotein.

steatosis may be associated with higher circulating leukocyte counts and markers for atherosclerosis, with more pronounced influences for peri-aortic adiposity. Our data suggested the differential biological impacts for region-specific visceral adiposity.

## Introduction

Excessive adipose tissue is associated with a chronic inflammatory status that leads to a variety of metabolic disorders including dyslipidemia, hypertension, and type 2 diabetes with region-specific properties [1]. Compared to subcutaneous adipose tissue, visceral adipose tissue is functionally more active in several metabolic derangements and may not be accurately identified by traditional anthropometrics including body weight and body mass index (BMI) assessment [2]. Over-production of various pro-inflammatory paracrine and mediators by visceral adipose tissue leads to obesity-related systemic inflammation and insulin resistance [3, 4]. On the other hand, fatty liver disease—characterized by excessive fatty infiltration of liver tissue—is an increasingly recognized cause of chronic liver disease worldwide, and is highly associated with central obesity, local hepatic inflammation, insulin resistance and increased systemic oxidative stress [5, 6].

Typically, the local inflammatory milieu of visceral adipose tissue is characterized by monocyte/macrophage infiltration and a diversity of lymphocyte subtypes [7, 8]. Additionally, increasing evidence suggests that various adipokines, free radicals from exaggerated oxidative stress and proinflammatory cytokines secreted directly from adipocytes may have remote adverse cardiovascular effects in obesity [9]. Although cytokines are rarely measured in daily practice, the total white blood cell count (WBC) and its subtypes (e.g. monocytes, lymphocytes, neutrophils, eosinophils, and basophils) may reflect a patient's inflammatory status in the absence of infection [10]. In particular, the neutrophil-to-lymphocyte ratio (NLR) has been proposed as a risk marker for adverse inflammatory status in metabolic syndrome, several cardiovascular disorders and cancer [11].

Recently, two volume-based measures of visceral adiposity burden—peri-cardial (PCF) and thoracic peri-aortic adipose tissue (TAT)—have been shown to correlate with metabolic risk profiles and atherosclerosis [12]. Studies over the past decade also suggest that both PCF and TAT are independently associated with systemic inflammatory markers such as hs-CRP. As region-specific visceral adiposity may have biologically diverse effects and associate with different pathological conditions [13, 14], the aim of this study was to investigate to what extent these visceral fat measures (including PCF, TAT and hepatic steatosis) may be associated with circulating leukocyte counts and the NLR in a large cohort of adults in Taiwan.

## Methods

### Study population

From 2005 to 2009, we studied consecutive subjects who underwent a comprehensive cardiovascular health survey at our center that included a non-contrast computed tomography (CT) scan of the chest to assess the presence and burden of coronary calcium. Subjects with CT data available comprised the main study population in our current work. This study complies with the Declaration of Helsinki, and the study protocol was approved by the Institutional Review Board at Mackay Memorial Hospital, Taipei, Taiwan. Data were analyzed anonymously.

Baseline demographics and medical history were obtained along with a detailed physical exam. Structured questionnaires were used to quantify self-reported alcohol consumption, smoking and physical activity. Presence of coronary arterial disease (CAD) was defined as a history of myocardial infarction or prior history of coronary revascularization. History of hypertension was defined as either systolic blood pressure higher than 140mmHg, diastolic pressure higher than 90mmHg or previous diagnosed hypertension on treatment. Diabetes was defined as current usage of any medications for diabetes or known diagnosis of diabetes. Hyperlipidemia was defined as known history of hyperlipidemia or medications (e.g. statin or fibrate). Anthropometric measures including height, weight, waist and hip circumferences were all obtained. Resting blood pressures were measured by medical staff using a standardized automated sphygmomanometer.


### Biochemical data, white blood cell counts and serum markers of systemic inflammation

Venous blood sampling and analysis were performed according to the Clinical Laboratory Standards Institute guidelines (Specimen Choice, Collection, and Handling; Approved Guideline H18-A3). Levels of total white blood cell (WBC) counts (leukocytes), neutrophils, lymphocytes, and monocytes were all determined by an automated blood cell counter utilizing Coulter LH780 Hematology Analyzer (Beckman Coulter Ireland Inc Mervue, Galway, Ireland). In our current work, absolute cell counts were used in the analyses. To ensure accuracy, results were verified by repeating the tests on the same tube one day later. A Hitachi 7170 Automatic Analyzer (Hitachi Corp. Hitachinaka Ibaraki, Japan) was used to measure fasting glucose, post-prandial glucose, HbA1c, uric acid, blood urea nitrogen, creatinine, homocysteine, and several lipid profiles including HDL, LDL, total cholesterol and triglyceride, with estimated glomerular filtration rate (eGFR) calculated using the Modification of Diet in Renal Disease equation. High-sensitivity CRP (hs-CRP) levels were determined using a highly sensitive, latex particle-enhanced immunoassay Elecsys 2010 (Roche, Mannheim, Germany).

### Measurements of ectopic visceral adipose tissue

Scans were performed using a 16-slice multidetector CT (MDCT) scanner (Sensation 16; Siemens Medical Solutions, Forchheim, Germany) with  $16 \times 0.75$  mm collimation, rotation time of 420 msec, and tube voltage of 120 kV. In one breath hold, images were acquired from above the level of tracheal bifurcation to below the base of heart using prospective electrocardiographic triggering, with the center of the acquisition at 70% of the R-R interval. From the raw data, the images were reconstructed with standard kernel in 3 mm thick axial, non-overlapping slices and 25 cm field of view.

PCF volumes were quantified from the heart CT scan using a dedicated workstation (Aquarius 3D Workstation, TeraRecon, San Mateo, CA, USA). The semi-automatic segmentation technique was developed for quantification of adipose tissue volumes. We traced pericardium in axial MDCT images manually from the level of left main coronary artery to diaphragm every four to six slices. The computer software then automatically interpolated and traced pericardium along the manually traced areas. All automatically traced slices were verified and modified if necessary for accuracy. Adipose tissue was defined as pixels within a window of -195 to -45 HU and a window center of -120 HU. Pericardial fat (PCF) was defined as any adipose tissue located within the pericardial sac. Thoracic peri-aortic adipose tissue (TAT) was defined as all of the adipose tissue surrounding the thoracic aorta extending 67.5 mm caudally from the level of the bifurcation of pulmonary arteries. (Fig 1) This approach has previously been validated [15, 16]. Two observers performed independent readings on a random



**Fig 1.** Examples of PCF (A, B) and TAT (C, D) in 2D and 3D computed tomography views. A 69-year-old male with large amounts of PCF (235.8ml), TAT (24.9ml) and the lab data (WBC: 10700/mm<sup>3</sup>, Neutrophil: 9502/mm<sup>3</sup>, NLR: 15.05, monocyte: 481/mm<sup>3</sup>) (First row). A 45-year-old female with small amounts of PCF (18.7ml), TAT (1.24ml) and the lab data (WBC: 3900/mm<sup>3</sup>, Neutrophil: 2211/mm<sup>3</sup>, NLR: 1.702, monocyte: 238/mm<sup>3</sup>) (Second row). PCF: peri-cardial adipose tissue; TAT: thoracic peri-aortic adipose tissue; WBC: white blood cell count; NLR: neutrophil-to-lymphocyte ratio.

<https://doi.org/10.1371/journal.pone.0207284.g001>

subset of 40 subjects. The intra-observer and inter-observer coefficient of variation were 4.27%, 4.87% and 6.58%, 6.81% for PCF and TAT, respectively [15].

### Ultrasonographic assessment on grade of hepatic steatosis

Hepatic ultrasonography was performed in all patients by experienced gastroenterologists, who were blinded to the patients' clinical data. Hepatic steatosis was diagnosed based on characteristic ultrasonographic characteristics, including diffuse hyper echogenicity of the liver relative to the kidneys, ultrasonography beam attenuation, and poor visualization of the intrahepatic vessel borders and diaphragm [17]. Ultrasonography allows detection of the presence of mild and moderate-to-severe hepatic steatosis as alternative surrogate of intra-abdominal adiposity, with a sensitivity and specificity of approximately 85% and 95%, respectively (when liver fat infiltration on histology is at least 20–30%)[17]. We categorized hepatic steatosis severity (none, mild, moderate, severe) based on the intensity of hepatic hyper echogenicity compared to the kidney.

### Statistical analysis

Baseline demographics were compared across quintiles of PCF and TAT volumes, and the Wilcoxon rank-sum test was used to estimate the statistical significance of trends across all ordered groups. The prevalence of hepatic steatosis according to PCF/TAT quintiles was visualized using bar graphs, and compared across categories using (chi-squared tests). The associations between leukocyte counts and PCF/TAT were explored using Pearson's correlation, and logistic regression was used to examine the association of leukocyte counts with moderate or severe hepatic steatosis. We further conducted uni- and multi-variate linear regression models

to explore the independent relationships of ectopic fat measures (PCF/TAT and hepatic steatosis as independent variables) with leukocyte counts (as dependent variables) after accounting for several key baseline clinical co-variables. Receiver operating characteristic curves with c-statistics were used to test the diagnostic performance of PCF/TAT and hepatic steatosis added to age, gender, BMI, biochemical profiles, lifestyle factors, and medical history to identify abnormally high total WBC counts ( $>6.9 \times 10^3/\mu\text{L}$ ) [18], and NLR ( $>2.51$ ) [19]. The potential confounding factors for multi-variate (MV) regression models included age, gender, BMI, blood pressure, fasting glucose, lipid profiles, renal function, medical histories of hypertension, diabetes, hyperlipidemia, coronary disease, smoking, and alcohol use.

All data were analyzed using STATA 12.0 (STATA Corp., College Station, Texas). All statistical tests were two-sided with  $p < 0.05$  considered to be statistically significant.

## Results

### Baseline characteristics and circulating individual white blood cell counts of study participants

A total of 3,291 subjects met our entry criteria and were enrolled in this study. The mean age was  $49.8 \pm 9.8$  years, with averaged BMI estimated to be  $24.7 \pm 3.5 \text{ kg/m}^2$  in our current cohort. Baseline demographic data across quintiles of PCF and TAT are displayed in Table 1, with mean ( $\pm$  standard deviation) PCF and TAT volumes of  $75.7 (\pm 30.9)$  and  $7.1 (\pm 4.0)$  ml, respectively. Among 3,196 subjects with liver coding data available, 2,736 (85.6%) had no or mild hepatic steatosis, 277 (8.7%) had moderate and 183 (5.7%) had severe hepatic steatosis.

Higher quintile of PCF and TAT volume was associated with more advanced age, higher systolic/diastolic blood pressures, increased body height and weight, greater BMI, higher fasting glucose, higher HbA1c, and unfavorable lipid profiles including higher total cholesterol, higher low-density lipoprotein (LDL), lower level of high-density lipoprotein (HDL), and lower estimated glomerular filtration rate (eGFR). Subjects with higher visceral fat depots were more likely to have prevalent hypertension, diabetes, hyperlipidemia and cardiovascular disease (all  $p < 0.05$ ).

### The association between visceral adipose tissue volume, hepatic steatosis and circulating differential white blood cell counts

Higher hs-CRP, total WBC and higher proportions of neutrophils, eosinophils, monocytes, lymphocytes, and myelocytes were observed with increasing amounts of PCF and TAT and with higher grades of liver steatosis (Table 2, all  $p < 0.01$ ).

Volumes of PCF/TAT and hepatic steatosis severity were positively correlated with hs-CRP, total WBC counts, and higher levels of neutrophils, eosinophils, monocytes, and lymphocytes (Table 3). In addition, subjects with higher quintile of PCF and TAT volume were associated with higher prevalence of hepatic steatosis (Fig 2). Volumes of PCF/TAT correlated with hs-CRP ( $r = 0.18, 0.21$  for PCF/TAT, both  $p < 0.001$ ). A graded increase of hs-CRP was also observed across hepatic steatosis categories (0.17, 0.19 & 0.25, trend  $p < 0.001$ ). Further, hs-CRP was positively associated with total WBC count ( $r = 0.24$ ) and several individual white blood cell components ( $r = 0.27, 0.16, 0.24$  for neutrophils, monocytes, and NLR, respectively, all  $p < 0.001$ ). Higher PCF and TAT along with substantially higher total WBC counts were observed across hepatic steatosis categories (Fig 3). In multi-variate models, adjustment for age, gender (Model 1), total body mass index (BMI, Model 2) and other clinical covariates attenuated the relationships between visceral adiposity (PCF/TAT), hepatic steatosis and hs-CRP, WBC or its differential blood cell types, especially for PCF and hepatic steatosis

**Table 1. Baseline characteristics of the study population by PCF and TAT quintiles.**

	PCF Quintiles (ml)						TAT Quintiles (ml)					
	Q1 (<50.1)	Q2 (50.1, 64.4)	Q3 (64.4, 77.6)	Q4 (77.6, 96.5)	Q5 (≥96.5)	p (trend)	Q1 (<3.8)	Q2 (3.8, 5.5)	Q3 (5.5, 7.4)	Q4 (7.4, 9.8)	Q5 (≥9.8)	p (trend)
<b>Baseline Characters</b>												
Age, year	45.14 ±9.28	48.72 ±8.88	49.88 ±9.13	51.04 ±9.50	54±9.95	<0.001	45.99 ±9.23	48.25±9.4	49.53 ±9.51	50.82 ±9.28	54.22 ±9.58	<0.001
Gender (male), (%)	350 (53.19%)	453 (68.84%)	502 (76.18%)	528 (80.24%)	548 (83.28%)	<0.001	193 (29.24%)	420 (63.83%)	539 (82.04%)	600 (90.91%)	629 (95.88%)	<0.001
Body height, cm	163.74 ±8.28	165.97 ±8.3	166.90 ±7.64	166.94 ±7.78	167.69 ±7.97	<0.001	161.64 ±8.02	165.68 ±8.58	167.17 ±8.03	168.34 ±6.65	168.4 ±7.16	<0.001
Body weight, kgw	59.48 ±10.09	65.15 ±9.97	68.73 ±10.53	71.58 ±11.51	76.33 ±12.13	<0.001	56.65 ±8.68	65.18 ±9.94	68.80 ±9.45	72.86 ±10.32	77.71 ±11.37	<0.001
BMI, kg/m <sup>2</sup>	22.08 ±2.71	23.59 ±2.73	24.59 ±2.89	25.59 ±3.15	27.07 ±3.57	<0.001	21.63 ±2.55	23.68 ±2.76	24.59 ±2.72	25.65 ±2.96	27.35±3.4	<0.001
SBP, mmHg	117.11 ±16.28	120.89 ±16.05	123.99 ±17.28	123.59 ±15.83	128.70 ±16.68	<0.001	115.00 ±16.04	120.19 ±16.34	122.94 ±15.26	125.48 ±15.49	130.68±17	<0.001
DBP, mmHg	71.49 ±10.40	74.88 ±10.79	77.02 ±10.9	77.21 ±10.18	79.63 ±10.21	<0.001	70.34 ±10.27	73.88 ±9.98	76.64 ±9.85	78.33 ±10.11	81.06 ±10.78	<0.001
Pulse rate, 1/min	72.17 ±10.20	71.76 ±10.82	73.36 ±9.63	73.1 ±11.01	73.95 ±12.03	0.002	72.53 ±10.21	71.55 ±11.18	72.96 ±10.01	73.09 ±10.73	74.18 ±11.6	<0.001
<b>Lab Data</b>												
Hb, g/dL	13.96 ±1.60	14.46 ±1.47	14.65 ±1.41	14.76 ±1.31	14.8±1.29	<0.001	13.45 ±1.51	14.33 ±1.53	14.73 ±1.23	15.02 ±1.15	15.1±1.13	<0.001
Fasting glucose, mg/dl	94.18 ±14.34	98.66 ±15.87	102.53 ±23.61	103.89 ±21.96	108.35 ±28.4	<0.001	92.88 ±9.87	98.33 ±17.47	101.38 ±21.61	102.72 ±19.9	112.45 ±30.91	<0.001
Total cholesterol, mg/dl	193.95 ±33.13	201.68 ±39.67	205.12 ±34.99	204.12 ±35.97	204.97 ±37.43	<0.001	196.24 ±35.23	201.44 ±36.03	205.19 ±36.41	204.41 ±38.8	202.51 ±35.36	<0.001
Triglyceride, mg/dl	103.7 ±64.04	129.83 ±168.89	143.64 ±88.22	148.42 ±89.52	169.44 ±114.91	<0.001	92.4 ±53.57	122.07 ±66.86	141.53 ±87.28	160.51 ±172.51	178.97 ±121.24	<0.001
LDL, cholesterol, mg/dl	120.75 ±29.56	130.10 ±32.03	134.13 ±31.61	134.06 ±32.76	134.32 ±33.25	<0.001	121.26 ±31.31	129.80 ±32.74	135.84 ±32.36	134.92 ±31.2	131.80 ±31.66	<0.001
HDL, cholesterol, mg/dl	59.80 ±15.81	54.68 ±14.27	51.92 ±13.03	50.17 ±11.97	47.28 ±11.84	<0.001	63.28 ±15.83	55.48 ±13.87	51.30 ±12.11	47.68 ±10.35	45.90 ±10.12	<0.001
sGPT, mg/dL	24.94 ±29.07	27.18 ±19.31	31.61 ±24.4	32.65 ±23.28	36.63 ±29.96	<0.001	22.01 ±25.94	27.36 ±23.16	30.85 ±19.57	35.83 ±31.99	36.98 ±23.55	<0.001
eGFR, ml/min/1.73m <sup>2</sup>	89.22 ±14.43	86.63 ±15.53	85.63 ±14.65	84.95 ±14.94	83.76 ±14.99	0.009	90.10 ±15.07	87.17 ±14.44	86.00 ±14.93	83.89 ±14.06	82.26 ±15.41	<0.001
<b>Medical History</b>												
History of hypertension, %	45(6.84%)	82 (12.46%)	116 (17.60%)	115 (17.48%)	176 (26.75%)	<0.001	38(5.76%)	70 (10.64%)	90(13.7%)	117 (17.73%)	219 (33.38%)	<0.001
History of diabetes, %	14(2.13%)	34(5.17%)	35(5.31%)	35(5.32%)	53(8.05%)	<0.001	11(1.67%)	23(3.5%)	32(4.87%)	39(5.91%)	66 (10.06%)	<0.001
History of hyperlipidemia treatment, %	13(1.98%)	30(4.56%)	27(4.1%)	38(5.78%)	56(8.51%)	<0.001	12(1.82%)	23(3.5%)	44(6.7%)	34(5.15%)	51(7.77%)	<0.001
History of CAD, %	12(1.82%)	19(2.89%)	23(3.49%)	36(5.47%)	52(7.9%)	<0.001	11(1.67%)	28(4.26%)	29(4.41%)	26(3.94%)	48(7.32%)	<0.001
Alcohol use, %	29(4.41%)	29(4.41%)	37(5.61%)	45(6.84%)	46(6.99%)	0.1	22(3.33%)	25(3.8%)	36(5.48%)	44(6.67%)	59(8.99%)	<0.001
Current smoker, %	50(7.6%)	55(8.36%)	72 (10.93%)	89 (13.53%)	96 (14.59%)	<0.001	30(4.55%)	53(8.05%)	76 (11.57%)	92 (13.94%)	111 (16.92%)	<0.001

BMI: body mass index; DBP: diastolic blood pressure; eGFR: estimated glomerular filtration rate; Hb: hemoglobin; HDL: high-density lipoprotein; LDL: low-density lipoprotein; SBP: systolic blood pressure; sGPT: serum glutamate-pyruvate transaminase.

<https://doi.org/10.1371/journal.pone.0207284.t001>

**Table 2. The correlation between regional-specific adipose tissue, WBC, various individual white blood cells and NLR by PCF, TAT quintiles and hepatic steatosis.**

A												
	PCF Quintiles (ml)						TAT Quintiles (ml)					
Factors	Q1 ( <b>&lt;50.1</b> )	Q2 ( <b>50.1, 64.4</b> )	Q3 ( <b>64.4, 77.6</b> )	Q4 ( <b>77.6, 96.5</b> )	Q5 ( <b>&gt;96.5</b> )	p (trend)	Q1 ( <b>&lt;3.8</b> )	Q2 ( <b>3.8, 5.5</b> )	Q3 ( <b>5.5, 7.4</b> )	Q4 ( <b>7.4, 9.8</b> )	Q5 ( <b>&gt;9.8</b> )	p (trend)
hs-CRP, mg/L	0.15 ±0.19	0.19±0.20	0.19±0.21	0.25±0.27	0.27 ±0.24	<0.001	0.15 ±0.19	0.17 ±0.18	0.21 ±0.22	0.25 ±0.25	0.27 ±0.27	<0.001
WBC count, (10 <sup>3</sup> /mm <sup>3</sup> )	5.72 ±1.39	5.98±1.34	6.07±1.33	6.25±1.38	6.37 ±1.34	<0.001	5.52 ±1.37	5.9±1.3	6.09 ±1.31	6.38 ±1.34	6.51 ±1.31	<0.001
Neutrophil count (10 <sup>3</sup> /mm <sup>3</sup> )	3.3±1.12	3.43±1.03	3.46±1.01	3.58±1.03	3.64 ±1.01	<0.001	3.19 ±1.11	3.33 ±1.01	3.47 ±0.98	3.68 ±1.03	3.73±1	<0.001
Eosinophil count (10 <sup>3</sup> /mm <sup>3</sup> )	0.16 ±0.18	0.15±0.13	0.15±0.14	0.16±0.14	0.18 ±0.16	<0.001	0.14 ±0.18	0.15 ±0.14	0.16 ±0.13	0.17 ±0.14	0.18 ±0.13	<0.001
Basophil count (10 <sup>3</sup> /mm <sup>3</sup> )	0.02 ±0.02	0.02±0.02	0.02±0.02	0.03±0.03	0.02 ±0.03	0.258	0.02 ±0.02	0.02 ±0.02	0.02 ±0.02	0.02 ±0.03	0.02 ±0.03	0.109
Monocyte count (10 <sup>3</sup> /mm <sup>3</sup> )	0.39 ±0.14	0.41±0.14	0.42±0.14	0.42±0.15	0.43 ±0.15	<0.001	0.37 ±0.14	0.4±0.14	0.41 ±0.15	0.43 ±0.15	0.45 ±0.14	<0.001
Lymphocyte count (10 <sup>3</sup> /mm <sup>3</sup> )	1.83 ±0.48	1.94±0.51	1.98±0.52	2.03±0.57	2.04 ±0.55	<0.001	1.78 ±0.48	1.94 ±0.49	1.99 ±0.54	2.04 ±0.53	2.08 ±0.56	<0.001
NLR	1.89 ±0.79	1.86±0.70	1.85±0.70	1.89±0.72	1.90 ±0.73	0.497	1.80 ±0.80	1.81 ±0.69	1.85 ±0.71	1.91 ±0.70	1.92 ±0.74	<0.001
B												
	Grade of Hepatic Steatosis											
Factors	No/Mild (n = 2736)		Moderate (n = 277)		Severe (n = 183)		p (trend)					
hs-CRP, mg/L	0.16±0.23		0.19±0.21		0.25±0.26		<0.001					
WBC count, (10 <sup>3</sup> /mm <sup>3</sup> )	5.99±1.57		6.28±1.55		6.74±1.70		<0.001					
Neutrophil count (10 <sup>3</sup> /mm <sup>3</sup> )	3.42±1.19		3.57±1.17		3.92±1.27		<0.001					
Eosinophil count (10 <sup>3</sup> /mm <sup>3</sup> )	0.16±0.15		0.16±0.14		0.19±0.15		0.001					
Basophil count (10 <sup>3</sup> /mm <sup>3</sup> )	0.02±0.02		0.02±0.02		0.02±0.03		0.69					
Monocyte count (10 <sup>3</sup> /mm <sup>3</sup> )	0.40±0.15		0.42±0.15		0.46±0.19		<0.001					
Lymphocyte count (10 <sup>3</sup> /mm <sup>3</sup> )	1.96±0.60		2.04±0.59		2.11±0.63		<0.001					
NLR	1.87±0.83		1.84±0.69		1.99±0.81		0.001					

WBC: white blood cell; NLR: neutrophil lymphocyte ratio

<https://doi.org/10.1371/journal.pone.0207284.t002>

(Table 4). By defining abnormal total WBC count and NLR at a cut-off of 5,900 x 10<sup>3</sup>μL and 2.51, respectively, higher PCF and TAT burden were significantly associated with higher risk of abnormally high WBC count (Crude OR: 1.25 [95% CI: 1.15–1.35] & 1.38 [95% CI: 1.28–1.49] for PCF & TAT, both p<0.001) and showed modest associations with abnormally high NLR (Crude OR: 1.08 [95% CI: 0.98–1.18] & 1.12 [95% CI: 1.02–1.23], p = 0.126 & 0.014), respectively. In fully adjusted models, only higher TAT was independently associated with abnormally high total WBC count and NLR (adj. OR: 1.18 [95% CI: 1.05–1.33] & 1.21 [95% CI: 1.05–1.39], p = 0.006 & 0.007 for total WBC and NLR, respectively) (Fig 4A and 4B). Existence of moderate/severe hepatic steatosis (vs. no or mild) was also associated with abnormally high total WBC count (Crude OR: 1.42 [95% CI: 1.24.15–1.64, P<0.001]) but not abnormal NLR (Crude OR: 1.00 [95% CI: 0.83.15–1.20, P = 0.98]; however, the association between hepatic steatosis and high WBC count was attenuated in fully adjusted models (adj. OR: 1.08, p = 0.34) (Fig 4A and 4B). Associations between TAT and total WBC, neutrophil, monocyte

**Table 3. The regression models for both visceral adipose tissue, hepatic steatosis (moderate-severe degree) with WBC, it's subtypes and NLR.**

<b>A</b>				
	PCF, ml		TAT, ml	
Factors	$\beta$ -Coef	p value	$\beta$ -Coef	p value
hs-CRP, mg/L	0.09	0.002	0.12	<0.001
WBC count ( $10^3/\text{mm}^3$ )	0.224	<0.001	0.32	<0.001
Neutrophil count ( $10^3/\text{mm}^3$ )	0.127	<0.001	0.197	<0.001
Eosinophil count ( $10^3/\text{mm}^3$ )	0.01	<0.001	0.015	<0.001
Basophil count ( $10^3/\text{mm}^3$ )	0	0.941	0	0.966
Monocyte count ( $10^3/\text{mm}^3$ )	0.015	<0.001	0.024	<0.001
Lymphocyte count ( $10^3/\text{mm}^3$ )	0.059	<0.001	0.078	<0.001
NLR	0.019	0.164	0.042	0.002
<b>B</b>				
	Hepatic Steatosis (Moderate-Severe Degree)			
Factors	Coef	p value		
hs-CRP, mg/L	0.07	0.003		
WBC count ( $10^3/\text{mm}^3$ )	0.35	<0.001		
Neutrophil count ( $10^3/\text{mm}^3$ )	0.23	<0.001		
Eosinophil count ( $10^3/\text{mm}^3$ )	0.014	0.007		
Basophil count ( $10^3/\text{mm}^3$ )	—	—		
Monocyte count ( $10^3/\text{mm}^3$ )	0.02	<0.001		
Lymphocyte count ( $10^3/\text{mm}^3$ )	0.077	<0.001		
NLR	0.03	0.01		

WBC: white blood cell; NLR: neutrophil lymphocyte ratio


<https://doi.org/10.1371/journal.pone.0207284.t003>

count, and NLR remained statistically significant in fully-adjusted models (Table 4, Model 3, all  $p < 0.05$ ). Fig 5A-5D demonstrate changes in the c-statistic for models of total WBC count and NLR as more co-variables were added: (1) age, gender, BMI; (2) model 1 + fasting sugar, LDL, HDL and eGFR; (3) model 2 + history of hypertension, diabetes, cardiovascular disease + lifestyle behaviors (active smoker and regular alcohol use); and (4) model 4 + PCF or TAT. In the final model 4, addition of TAT—but not PCF—statistically significantly increased the C-statistic for both total WBC count and NLR (Fig 5A-5D).

## Discussion

In this large cohort of subjects who underwent a cardiovascular health survey in Taiwan, we demonstrate that 3 measures of visceral adiposity—PCF, TAT and hepatic steatosis—are associated with circulating WBC counts and the systemic inflammatory marker hs-CRP. Moreover, compared to PCF, TAT appears to be more strongly related to WBC counts and the NLR, an integrative measure of generalized inflammation, even after accounting for clinical co-variables. Further, we observed that TAT added independent and incremental value in identifying abnormally higher total WBC counts and NLR. Our study suggests that regional ectopic fat deposition of the thoracic aorta (TAT) is associated with systemic inflammation and altered leukocyte numbers above and beyond clinical characteristics and body anthropometrics.

Peripheral total leukocyte counts (WBC) and its subtypes have previously been associated with cardiometabolic risk factors including dyslipidemia, obesity, and metabolic syndrome [20–22]. Higher total WBC counts have been further associated with clinical events such as incident type 2 diabetes and increased mortality after acute coronary syndromes [23, 24].




**Fig 2. Prevalent hepatic steatosis across PCF, TAT volume quintiles.** Subjects with higher quintile of PCF and TAT volume were more likely to have prevalent Hepatic Steatosis (both  $p < 0.001$  for  $\chi^2$  test: 32.2 & 38.7, respectively).

<https://doi.org/10.1371/journal.pone.0207284.g002>

These and other relevant studies illustrate how atherosclerotic cardiovascular disease is a complex multi-factorial process in which inflammation plays a major role. Abnormal adipose tissue distribution and function are believed to contribute to atherosclerosis, an effect which may be mediated by inflammation. At the adipose tissue level, inflammation is complex and characterized by dynamic populations of infiltrating neutrophils, macrophages, mast cells, and lymphocytes [7, 25]. Among these, adipose tissue macrophages (ATMs), may be the most important drivers of obesity-related chronic inflammation [26]. In both mouse models and human subjects, adipocyte hypertrophy induced by excessive caloric intake triggers macrophage accumulation in visceral adipose depots where they form so-called “crown-like structures” (CLSs) around dead adipocytes [27, 28]. These infiltrating macrophages express the pro-inflammatory M1 phenotype and secrete a variety of inflammatory cytokines such as tumor necrosis factor- $\alpha$ , interleukin-6 and monocyte chemoattractant protein-1, leading to up-regulated circulating leukocytes and trigger systemic inflammatory responses [29]. Importantly, these pro-inflammatory cytokines may act locally on adipocyte function and systemically on other tissues to promote the pathogenesis of cardiometabolic disease [30]. In our current work, we demonstrate that both increased PCF and TAT (but not hepatic steatosis) were independently associated with higher total leukocyte counts after accounting for clinical co-variables. Of note, only TAT showed independent and incremental value in predicting abnormally high total leukocyte counts ( $5900 \times 10^3/\mu\text{L}$ ) and higher NLR ( $>2.51$ ).

Recently, the NLR has been proposed as a simple, clinically available and inexpensive marker of systemic inflammation, which is helpful for risk stratification of patients with cardiovascular diseases, diabetes mellitus, metabolic syndrome [11, 31], and for subsequent adverse cardiovascular events following myocardial infarction [32]. Neutrophils may contribute to acute myocardial injury by degranulation and secretion of inflammatory cytokines and proteases, resulting in tissue damage and reperfusion injury [33, 34]. Furthermore, neutrophils can also invade atherosclerotic plaques and trigger oxidative stress by releasing superoxide



**Fig 3. PCF, TAT, NLR, total leukocyte counts across hepatic steatosis category.** Higher severity of Hepatic Steatosis [categorized as (1) No/Mild, (2) Moderate, and (3) Severe] was associated with greater burden of visceral adiposity (A and B) and higher total leukocyte counts (D), but was not associated with NLR (C). \* $p < 0.05$  compared to Non- Hepatic Steatosis group, † $p < 0.05$  compared to Moderate Hepatic Steatosis group.

<https://doi.org/10.1371/journal.pone.0207284.g003>

radicals, proteolytic enzymes & arachidonic acid derivatives, which all make atherosclerotic plaques more vulnerable [35]. On the other hand, lymphocytes play an important role in regulating the inflammatory response and are suppressed by corticosteroids in response to the stress of acute coronary syndrome [36]. In previous studies, however, increased BMI was associated with total WBC and individual sub-types but not with NLR [37–39]. A multi-center study revealed that NLR is positively correlated with hepatic steatosis activity score, pro-inflammatory cytokines and CRP [40], though in our multi-variate analysis the associations between hepatic steatosis and all types of leucocytes were attenuated, indicating that hepatic steatosis might play a lesser role in mediating systemic leucocytes circulation. In aggregate, our work further expands on these previous observations to show that TAT—a measure of regional visceral adiposity—, instead of PCF or hepatic steatosis, is tightly associated with NLR beyond clinical covariates, BMI and lipid profiles. These findings are also consistent with ectopic visceral fat contributing to elevated cardiovascular risk in “metabolically obese” normal-weight individuals [41, 42]. While both TAT and PCF remained independently related to total leukocyte counts and NLR, only TAT statistically significantly improved the c-statistic of our fully-

**Table 4.** Multivariate adjustment models in the association between both visceral adipose tissue, moderate-to-severe hepatic steatosis, WBC, it's subtypes and NLR.

Multi-variate Model	Multi-variate Model 1				Multi-variate Model 2				Multi-variate Model 3			
Factor	PCF, ml		TAT, ml		PCF,ml		TAT, ml		PCF, ml		TAT, ml	
	β-Coef	p value	β-Coef	p value	β-Coef	p value	β-Coef	p value	β-Coef	p value	β-Coef	p value
hs-CRP, mg/L	0.06	0.032	0.1	0.001	0.06	0.23	0.06	0.094	0.05	0.52	0.08	0.38
WBC count ( $10^3/\text{mm}^3$ )	0.243	<0.001	0.339	<0.001	0.107	<0.001	0.208	<0.001	0.135	0.012	0.155	0.011
Neutrophil count ( $10^3/\text{mm}^3$ )	0.15	<0.001	0.242	<0.001	0.067	0.004	0.169	<0.001	0.065	0.143	0.106	0.04
Eosinophil count ( $10^3/\text{mm}^3$ )	0.008	0.005	0.01	0.002	0.006	0.099	0.007	0.056	0.004	0.42	0.157	0.073
Basophil count ( $10^3/\text{mm}^3$ )	0	0.774	0	0.537	0	0.945	0	0.635	0	0.718	0.001	0.27
Monocyte count ( $10^3/\text{mm}^3$ )	0.015	<0.001	0.021	<0.001	0.007	0.026	0.013	<0.001	0.007	0.243	0.017	0.022
Lymphocyte count ( $10^3/\text{mm}^3$ )	0.059	<0.001	0.068	<0.001	0.023	0.054	0.023	0.089	0.033	0.12	0.028	0.248
NLR	0.028	0.052	0.072	<0.001	0.02	0.218	0.081	<0.001	0.011	0.52	0.062	0.002


  

Multi-variate Model	(Moderate-to-Severe) Hepatic Steatosis					
Factor	Multi-variate Model 1		Multi-variate Model 2		Multi-variate Model 3	
	Coef	p value	Coef	p value	Coef	p value
hs-CRP, mg/L	0.07	0.004	0.01	0.56	0.004	0.71
WBC count ( $10^3/\text{mm}^3$ )	0.28	<0.001	0.13	0.01	—	—
Neutrophil count ( $10^3/\text{mm}^3$ )	0.19	<0.001	0.1	0.012	—	—
Eosinophil count ( $10^3/\text{mm}^3$ )	—	—	—	—	—	—
Basophil count ( $10^3/\text{mm}^3$ )	—	—	—	—	—	—
Monocyte count ( $10^3/\text{mm}^3$ )	0.02	0.001	—	—	—	—
Lymphocyte count ( $10^3/\text{mm}^3$ )	0.06	0.007	—	—	—	—
NLR	0.04	0.014	0.03	0.032	—	—

Model 1: adjusted for age, gender; Model 2: adjusted for age, gender, BMI; Model 3: adjusted for age, gender, BMI, multi-variate models (MV); BMI: body mass index. MV: systolic blood pressure, pulse rate, fasting glucose, HDL, LDL, eGFR, medical history of diabetes, hypertension, cardiovascular disease, current smoker, alcohol use.


<https://doi.org/10.1371/journal.pone.0207284.t004>

adjusted model. This finding may be partly explained by anatomical difference. Elicited abdominal visceral adipose tissue inflammation close to portal circulation may release fatty acids and cytokines that directly affect liver (the end-organ), which is considered as the key factor of hepatic steatosis and insulin resistance [43]. Instead, PCF is anatomically confined



**Fig 4.** Odds of abnormally high WBC (a) and NLR (b) in relation to increasing PCF, TAT, and presence of hepatic steatosis. The crude and adjusted (adj.) odds ratios (OR) for PCF, TAT and Hepatic Steatosis in identifying abnormally high total leukocyte counts (WBC, cut-off:  $>6.9 \times 10^3/\mu\text{L}$ ) and NLR (cut-off:  $>2.51$ ).

<https://doi.org/10.1371/journal.pone.0207284.g004>



**Fig 5. The prediction value of abnormally high WBC, NLR by PCF, TAT.** The C-statistics of PCF, and TAT in the model prediction for abnormally high total leukocyte counts (WBC, cut-off:  $>6.9 \times 10^3/\mu\text{L}$ )<sup>18</sup> and NLR (cut-off:  $>2.51$ ). Again, TAT demonstrated significant incremental value beyond conventional cardiovascular risk factors by C-statistics from 0.66 to 0.67 for abnormal WBC, and 0.50 to 0.54 for abnormal NLR (c, d). # $p < 0.05$  in fully adjusted models.

<https://doi.org/10.1371/journal.pone.0207284.g005>

within the pericardial sac, which was believed to be the source of pro-inflammatory adipokines near coronary arteries and myocardium. Recent studies have demonstrated close relationships between PCF and pathogenesis of atherosclerosis or certain degree of myocardial dysfunction [44, 45]. Compared to PCF and hepatic steatosis, TAT may play a more important role in systemic inflammatory process, partly due to the fact that cytokines and pro-inflammatory mediators from TAT can more easily diffuse and distribute through the adventitia layer across the arterial wall to enter the systemic circulation [12, 46]. This data together with our previous findings underscores the possibility of using TAT as important metabolic surrogate marker which may provide additional information in identifying subjects at higher risk of cardiovascular disorders. Further investigation is warranted to explain why different locations of ectopic visceral fat may be differentially associated with systemic inflammation.

This study has some limitations. Because it is a cross-sectional study, causal inferences cannot be made and the possibility of residual or unknown confounding cannot be excluded. Additionally, our subjects may not represent the general population in Taiwan, since they were participating in a voluntary health survey and were not randomly selected. Our study was disproportionately male, and so further study in larger female populations is warranted. Finally, our study represented only a single ethnicity (Taiwanese). In light of racial and ethnic differences in NLR [47], future studies should examine these relationships in other populations.

## Conclusion

Our study results indicate that greater visceral fat depots appear to be associated with higher WBC counts and individual cell types independent of body mass index and other traditional risk factors; however, compared to ectopic fat within the pericardial sac, visceral fat surrounding the aorta is more strongly associated with the NLR.

## Supporting information

**S1 Fig. The relationship between total leukocyte, neutrophil, monocyte, NLR and groups according to low or high BMI ( $<24.3$ ,  $\geq 24.3$  kg/m<sup>2</sup>) and low and high PCF/TAT categories ( $<71$ ,  $\geq 71$  ml for PCF;  $<6.4$ ,  $\geq 6.4$  ml for TAT) based on median values.** Total WBC and the proportion of neutrophil and monocyte tended to increase across BMI and both PCF and TAT groups (all  $p$  for trend:  $<0.05$ ) For NLR, the association across BMI/TAT categories was statistically significant but was not for BMI/PCF categories ( $p = 0.008$  vs.  $p = 0.196$  for TAT vs. PCF, respectively). (PPTX)

## Author Contributions

**Data curation:** Richard Kuo, Ta-Chuan Hung, Chuan-Chuan Liu, Charles Jia-Yin Hou, David A. Zidar.

**Formal analysis:** Charles Jia-Yin Hou.

**Investigation:** Shun-Chuan Chang.

**Methodology:** Chuan-Chuan Liu.

**Validation:** Chung-Lieh Hung, Hung-I Yeh.

**Visualization:** Jing-Yi Sun, Tung-Hsin Wu.

**Writing – original draft:** Kuo-Tzu Sung, Chun-Ho Yun.

**Writing – review & editing:** Chun-Ho Yun, Chung-Lieh Hung, Ricardo C. Cury, David A. Zidar, Hiram G. Bezerra, Chris T. Longenecker.

## References

1. Glass CK, Olefsky JM. Inflammation and lipid signaling in the etiology of insulin resistance. *Cell Metab.* 2012; 15(5):635–45. <https://doi.org/10.1016/j.cmet.2012.04.001> PMID: 22560216
2. Bays HE, Gonzalez-Campoy JM, Bray GA, Kitabchi AE, Bergman DA, Schorr AB, et al. Pathogenic potential of adipose tissue and metabolic consequences of adipocyte hypertrophy and increased visceral adiposity. *Expert Rev Cardiovasc Ther.* 2008; 6(3):343–68. <https://doi.org/10.1586/14779072.6.3.343> PMID: 18327995
3. Reyes M, Gahagan S, Diaz E, Blanco E, Leiva L, Lera L, et al. Relationship of adiposity and insulin resistance mediated by inflammation in a group of overweight and obese Chilean adolescents. *Nutr J.* 2011; 10:4. <https://doi.org/10.1186/1475-2891-10-4> PMID: 21235793
4. Marques-Vidal P, Bastardot F, von Kanel R, Paccaud F, Preisig M, Waeber G, et al. Association between circulating cytokine levels, diabetes and insulin resistance in a population-based sample (CoLaus study). *Clin Endocrinol (Oxf).* 2013; 78(2):232–41. <https://doi.org/10.1111/j.1365-2265.2012.04384.x> PMID: 22409372
5. Videla LA, Rodrigo R, Orellana M, Fernandez V, Tapia G, Quinones L, et al. Oxidative stress-related parameters in the liver of non-alcoholic fatty liver disease patients. *Clin Sci (Lond).* 2004; 106(3):261–8. <https://doi.org/10.1042/CS20030285> PMID: 14556645

6. Marchesini G, Brizi M, Morselli-Labate AM, Bianchi G, Bugianesi E, McCullough AJ, et al. Association of nonalcoholic fatty liver disease with insulin resistance. *Am J Med.* 1999; 107(5):450–5. [https://doi.org/10.1016/S0002-9343\(99\)00271-5](https://doi.org/10.1016/S0002-9343(99)00271-5) PMID: 10569299
7. Liu J, Divoux A, Sun J, Zhang J, Clement K, Glickman JN, et al. Genetic deficiency and pharmacological stabilization of mast cells reduce diet-induced obesity and diabetes in mice. *Nat Med.* 2009; 15(8):940–5. <https://doi.org/10.1038/nm.1994> PMID: 19633655
8. Lolmede K, Duffaut C, Zakaroff-Girard A, Bouloumie A. Immune cells in adipose tissue: key players in metabolic disorders. *Diabetes Metab.* 2011; 37(4):283–90. <https://doi.org/10.1016/j.diabet.2011.03.002> PMID: 21507694
9. Bagi Z, Broskova Z, Feher A. Obesity and coronary microvascular disease—implications for adipose tissue-mediated remote inflammatory response. *Curr Vasc Pharmacol.* 2014; 12(3):453–61. PMID: 24846234
10. Stock W, Hoffman R. White blood cells 1: non-malignant disorders. *Lancet.* 2000; 355(9212):1351–7. [https://doi.org/10.1016/S0140-6736\(00\)02125-5](https://doi.org/10.1016/S0140-6736(00)02125-5) PMID: 10776761
11. Wang X, Zhang G, Jiang X, Zhu H, Lu Z, Xu L. Neutrophil to lymphocyte ratio in relation to risk of all-cause mortality and cardiovascular events among patients undergoing angiography or cardiac revascularization: a meta-analysis of observational studies. *Atherosclerosis.* 2014; 234(1):206–13. <https://doi.org/10.1016/j.atherosclerosis.2014.03.003> PMID: 24681815
12. Spiroglou SG, Kostopoulos CG, Varakis JN, Papadaki HH. Adipokines in periaortic and epicardial adipose tissue: differential expression and relation to atherosclerosis. *J Atheroscler Thromb.* 2010; 17(2):115–30. <https://doi.org/10.5551/jat.1735> PMID: 20145358
13. Yun CH, Bezerra HG, Wu TH, Yang FS, Liu CC, Wu YJ, et al. The normal limits, subclinical significance, related metabolic derangements and distinct biological effects of body site-specific adiposity in relatively healthy population. *PLoS One.* 2013; 8(4):e61997. <https://doi.org/10.1371/journal.pone.0061997> PMID: 23620798
14. Yun CH, Longenecker CT, Chang HR, Mok GS, Sun JY, Liu CC, et al. The association among peri-aortic root adipose tissue, metabolic derangements and burden of atherosclerosis in asymptomatic population. *J Cardiovasc Comput Tomogr.* 2016; 10(1):44–51. <https://doi.org/10.1016/j.jcct.2015.10.002> PMID: 26507645
15. Yun CH, Lin TY, Wu YJ, Liu CC, Kuo JY, Yeh HI, et al. Pericardial and thoracic peri-aortic adipose tissues contribute to systemic inflammation and calcified coronary atherosclerosis independent of body fat composition, anthropometric measures and traditional cardiovascular risks. *Eur J Radiol.* 2012; 81(4):749–56. <https://doi.org/10.1016/j.ejrad.2011.01.035> PMID: 21334840
16. Mahabadi AA, Massaro JM, Rosito GA, Levy D, Murabito JM, Wolf PA, et al. Association of pericardial fat, intrathoracic fat, and visceral abdominal fat with cardiovascular disease burden: the Framingham Heart Study. *Eur Heart J.* 2009; 30(7):850–6. <https://doi.org/10.1093/eurheartj/ehn573> PMID: 19136488
17. Hernaez R, Lazo M, Bonekamp S, Kamel I, Brancati FL, Guallar E, et al. Diagnostic accuracy and reliability of ultrasonography for the detection of fatty liver: a meta-analysis. *Hepatology.* 2011; 54(3):1082–90. <https://doi.org/10.1002/hep.24452> PMID: 21618575
18. Samdahl E, Bergstrom I, Brodin VP, Nijm J, Lundqvist Setterud H, Jonasson L. Neutrophil activation status in stable coronary artery disease. *PloS one.* 2007; 2(10):e1056. <https://doi.org/10.1371/journal.pone.0001056> PMID: 17957240
19. Kaya A, Kurt M, Tanboga IH, Isik T, Gunaydin ZY, Kaya Y, et al. Relation of neutrophil to lymphocyte ratio with the presence and severity of stable coronary artery disease. *Clinical and applied thrombosis/hemostasis: official journal of the International Academy of Clinical and Applied Thrombosis/Hemostasis.* 2014; 20(5):473–7. <https://doi.org/10.1177/1076029612473517> PMID: 23344996
20. Lee YJ, Lee HR, Shim JY, Moon BS, Lee JH, Kim JK. Relationship between white blood cell count and nonalcoholic fatty liver disease. *Dig Liver Dis.* 2010; 42(12):888–94. <https://doi.org/10.1016/j.dld.2010.04.005> PMID: 20472517
21. Oda E, Kawai R, Aizawa Y. Lymphocyte count was significantly associated with hyper-LDL cholesterolemia independently of high-sensitivity C-reactive protein in apparently healthy Japanese. *Heart Vessels.* 2012; 27(4):377–83. <https://doi.org/10.1007/s00380-011-0157-x> PMID: 21655904
22. Shim WS, Kim HJ, Kang ES, Ahn CW, Lim SK, Lee HC, et al. The association of total and differential white blood cell count with metabolic syndrome in type 2 diabetic patients. *Diabetes Res Clin Pract.* 2006; 73(3):284–91. <https://doi.org/10.1016/j.diabetes.2006.02.001> PMID: 16563549
23. Vozarova B, Weyer C, Lindsay RS, Pratley RE, Bogardus C, Tataranni PA. High white blood cell count is associated with a worsening of insulin sensitivity and predicts the development of type 2 diabetes. *Diabetes.* 2002; 51(2):455–61. <https://doi.org/10.2337/diabetes.51.2.455> PMID: 11812755

24. Barron HV, Cannon CP, Murphy SA, Braunwald E, Gibson CM. Association between white blood cell count, epicardial blood flow, myocardial perfusion, and clinical outcomes in the setting of acute myocardial infarction: a thrombolysis in myocardial infarction 10 substudy. *Circulation*. 2000; 102(19):2329–34. <https://doi.org/10.1161/01.CIR.102.19.2329> PMID: 11067784
25. Nishimura S, Manabe I, Nagasaki M, Eto K, Yamashita H, Ohsugi M, et al. CD8+ effector T cells contribute to macrophage recruitment and adipose tissue inflammation in obesity. *Nat Med*. 2009; 15(8):914–20. <https://doi.org/10.1038/nm.1964> PMID: 19633658
26. Wentworth JM, Naselli G, Brown WA, Doyle L, Phipson B, Smyth GK, et al. Pro-inflammatory CD11c +CD206+ adipose tissue macrophages are associated with insulin resistance in human obesity. *Diabetes*. 2010; 59(7):1648–56. <https://doi.org/10.2337/db09-0287> PMID: 20357360
27. Weisberg SP, McCann D, Desai M, Rosenbaum M, Leibel RL, Ferrante AW Jr., Obesity is associated with macrophage accumulation in adipose tissue. *The Journal of clinical investigation*. 2003; 112(12):1796–808. <https://doi.org/10.1172/JCI19246> PMID: 14679176
28. Xu H, Barnes GT, Yang Q, Tan G, Yang D, Chou CJ, et al. Chronic inflammation in fat plays a crucial role in the development of obesity-related insulin resistance. *J Clin Invest*. 2003; 112(12):1821–30. <https://doi.org/10.1172/JCI19451> PMID: 14679177
29. Lumeng CN, Bodzin JL, Saltiel AR. Obesity induces a phenotypic switch in adipose tissue macrophage polarization. *J Clin Invest*. 2007; 117(1):175–84. <https://doi.org/10.1172/JCI29881> PMID: 17200717
30. Hotamisligil GS, Shargill NS, Spiegelman BM. Adipose expression of tumor necrosis factor- $\alpha$ : direct role in obesity-linked insulin resistance. *Science*. 1993; 259(5091):87–91. <https://doi.org/10.1126/science.7678183> PMID: 7678183
31. Buyukkaya E, Karakas MF, Karakas E, Akcay AB, Tanboga IH, Kurt M, et al. Correlation of neutrophil to lymphocyte ratio with the presence and severity of metabolic syndrome. *Clin Appl Thromb Hemost*. 2014; 20(2):159–63. <https://doi.org/10.1177/1076029612459675> PMID: 22992349
32. Kaya MG, Akpek M, Lam YY, Yarlioglu M, Celik T, Gunebakmaz O, et al. Prognostic value of neutrophil/lymphocyte ratio in patients with ST-elevated myocardial infarction undergoing primary coronary intervention: a prospective, multicenter study. *Int J Cardiol*. 2013; 168(2):1154–9. <https://doi.org/10.1016/j.ijcard.2012.11.074> PMID: 23219132
33. Tamhane UU, Aneja S, Montgomery D, Rogers EK, Eagle KA, Gurm HS. Association between admission neutrophil to lymphocyte ratio and outcomes in patients with acute coronary syndrome. *The American journal of cardiology*. 2008; 102(6):653–7. <https://doi.org/10.1016/j.amjcard.2008.05.006> PMID: 18773982
34. Maxwell SR, Lip GY. Reperfusion injury: a review of the pathophysiology, clinical manifestations and therapeutic options. *Int J Cardiol*. 1997; 58(2):95–117. [https://doi.org/10.1016/S0167-5273\(96\)02854-9](https://doi.org/10.1016/S0167-5273(96)02854-9) PMID: 9049675
35. Eriksson EE, Xie X, Werr J, Thoren P, Lindbom L. Direct viewing of atherosclerosis in vivo: plaque invasion by leukocytes is initiated by the endothelial selectins. *Faseb j*. 2001; 15(7):1149–57. <https://doi.org/10.1096/fj.00-0537com> PMID: 11344083
36. Onsrud M, Thorsby E. Influence of in vivo hydrocortisone on some human blood lymphocyte subpopulations. I. Effect on natural killer cell activity. *Scand J Immunol*. 1981; 13(6):573–9. <https://doi.org/10.1111/j.1365-3083.1981.tb00171.x> PMID: 7313552
37. Balta S, Kurtoglu E, Kucuk U, Demirkol S, Ozturk C. Neutrophil-lymphocyte ratio as an important assessment tool. *Expert Rev Cardiovasc Ther*. 2014; 12(5):537–8. <https://doi.org/10.1586/14779072.2014.902309> PMID: 24745892
38. Furuncuoglu Y, Tulgar S, Dogan AN, Cakar S, Tulgar YK, Cakiroglu B. How obesity affects the neutrophil/lymphocyte and platelet/lymphocyte ratio, systemic immune-inflammatory index and platelet indices: a retrospective study. *Eur Rev Med Pharmacol Sci*. 2016; 20(7):1300–6. PMID: 27097950
39. Ryder E, Diez-Ewald M, Mosquera J, Fernandez E, Pedraza A, Vargas R, et al. Association of obesity with leukocyte count in obese individuals without metabolic syndrome. *Diabetes Metab Syndr*. 2014; 8(4):197–204. <https://doi.org/10.1016/j.dsx.2014.09.002> PMID: 25301008
40. Abdel-Razik A, Mousa N, Shabana W, Refaey M, ElMahdy Y, Elhelaly R, et al. A novel model using mean platelet volume and neutrophil to lymphocyte ratio as a marker of nonalcoholic steatohepatitis in NAFLD patients: multicentric study. *Eur J Gastroenterol Hepatol*. 2016; 28(1):e1–9. <https://doi.org/10.1097/MEG.0000000000000486> PMID: 26469357
41. Despres JP. Excess visceral adipose tissue/ectopic fat the missing link in the obesity paradox? *J Am Coll Cardiol*. 2011; 57(19):1887–9. <https://doi.org/10.1016/j.jacc.2010.10.063> PMID: 21545945
42. Chandra A, Neeland IJ, Berry JD, Ayers CR, Rohatgi A, Das SR, et al. The relationship of body mass and fat distribution with incident hypertension: observations from the Dallas Heart Study. *J Am Coll Cardiol*. 2014; 64(10):997–1002. <https://doi.org/10.1016/j.jacc.2014.05.057> PMID: 25190234

43. Rytka JM, Wueest S, Schoenle EJ, Konrad D. The portal theory supported by venous drainage-selective fat transplantation. *Diabetes*. 2011; 60(1):56–63. <https://doi.org/10.2337/db10-0697> PMID: [20956499](https://pubmed.ncbi.nlm.nih.gov/20956499/)
44. van Woerden G, Gorter TM, Westenbrink BD, Willems TP, van Veldhuisen DJ, Rienstra M. Epicardial fat in heart failure patients with mid-range and preserved ejection fraction. *Eur J Heart Fail*. 2018. <https://doi.org/10.1002/ejhf.1283> PMID: [30070041](https://pubmed.ncbi.nlm.nih.gov/30070041/)
45. Yorgun H, Canpolat U, Aytemir K, Hazirolan T, Sahiner L, Kaya EB, et al. Association of epicardial and peri-atrial adiposity with the presence and severity of non-valvular atrial fibrillation. *Int J Cardiovasc Imaging*. 2015; 31(3):649–57. <https://doi.org/10.1007/s10554-014-0579-5> PMID: [25466809](https://pubmed.ncbi.nlm.nih.gov/25466809/)
46. Tsuruda T, Kato J, Hatakeyama K, Kojima K, Yano M, Yano Y, et al. Adventitial mast cells contribute to pathogenesis in the progression of abdominal aortic aneurysm. *Circ Res*. 2008; 102(11):1368–77. <https://doi.org/10.1161/CIRCRESAHA.108.173682> PMID: [18451339](https://pubmed.ncbi.nlm.nih.gov/18451339/)
47. Azab B, Camacho-Rivera M, Taioli E. Average values and racial differences of neutrophil lymphocyte ratio among a nationally representative sample of United States subjects. *PLoS One*. 2014; 9(11): e112361. <https://doi.org/10.1371/journal.pone.0112361> PMID: [25375150](https://pubmed.ncbi.nlm.nih.gov/25375150/)