

Baptist Health South Florida

Scholarly Commons @ Baptist Health South Florida

All Publications

4-2019

Maximum IV tPA Dose for Obese Patients is Associated with Greater Likelihood of Hemorrhagic Conversion and Worse Functional Outcome at Discharge

Amy Starosciak

Baptist Health South Florida, amyst@baptisthealth.net

Felipe De Los Rios La Rosa

Baptist Hospital of Miami; Baptist Neuroscience Center, felipedl@baptisthealth.net

Follow this and additional works at: <https://scholarlycommons.baptisthealth.net/se-all-publications>



Part of the [Cardiology Commons](#), [Cardiovascular Diseases Commons](#), [Nervous System Diseases Commons](#), and the [Neurology Commons](#)

Citation

Starosciak, Amy and De Los Rios La Rosa, Felipe, "Maximum IV tPA Dose for Obese Patients is Associated with Greater Likelihood of Hemorrhagic Conversion and Worse Functional Outcome at Discharge" (2019). *All Publications*. 3132.

<https://scholarlycommons.baptisthealth.net/se-all-publications/3132>

This Conference Poster -- Open Access is brought to you for free and open access by Scholarly Commons @ Baptist Health South Florida. It has been accepted for inclusion in All Publications by an authorized administrator of Scholarly Commons @ Baptist Health South Florida. For more information, please contact Carrief@baptisthealth.net.

INTRODUCTION

IV tissue plasminogen activator (tPA, alteplase) is the mainstay of treatment for acute ischemic stroke at a weight-based dose of 0.9 mg/kg with a maximum limit of 90 mg for a person >100 kg (220 lb) according to United States (US) guidelines. The prevalence of adult obesity in the US has progressively increased; hence, the percentage of patients receiving the maximum dose is expected to rise as well. **We examined differences in patient characteristics and outcomes in acute ischemic stroke (AIS) patients who were treated with the weight-based dose (WBD) vs. the maximum dose (MD) of IV tPA.**

METHODS

We performed a historical cohort study using the local Get With The Guidelines-Stroke database from October 2013 to April 2017. Selection criteria included hospital admission, age ≥ 18 years, received IV tPA as treatment for AIS, and had a recorded weight (Figure 1).




Figure 1. The patient population selection based on exclusion and inclusion criteria

Patients were dichotomized into **WBD group** (<90 mg), weighing <100 kg, and **MD group** (90 mg), weighing ≥ 100 kg. We analyzed categorical variables using Chi square tests and continuous variables using independent samples t-tests.

RESULTS

Characteristics	Weight-Based Dose (n=290)	Max Dose (n=38)	p-Value
	Mean (range)	Mean (range)	
Age, mean (range)	77 (24-104)	62 (19-98)	<0.001
Weight (kg)	71 (32-99)	116 (100-182)	<0.001
	n (%)	n (%)	
Gender			
Male	119 (41)	29 (76)	<0.001
Female	171 (59)	9 (24)	
Ethnicity			
Hispanic	193 (67)	19 (50)	0.045
Non-Hispanic	97 (33)	19 (50)	
Initial NIHSS			
Low (0-4)	39 (13)	9 (24)	0.241
Moderate (5-10)	108 (37)	12 (32)	
High (>10)	143 (50)	17 (44)	
Disposition			
Good (Home/rehab)	201 (69)	24 (63)	0.442
Poor	89 (31)	14 (37)	
Discharge mRS			
0-2	66 (23)	7 (18)	<0.001
3-6	224 (77)	31 (82)	
90 day mRS			
0-2	100 (34)	9 (24)	0.627
3-6	111 (57)	14 (37)	
Lost to follow-up	79 (27)	11 (29)	
Complications			
None Serious	279 (97)	37 (88)	0.038
Pneumonia	16 (6)	3 (7)	0.671
Serious Systemic Hemorrhage	2 (1)	1 (2)	0.33
Symptomatic ICH	9 (3)	4 (10)	0.045

Table 1. Summary of patient characteristics and outcomes

- Patients in the **MD group** tended to be younger, male, non-Hispanic, and more likely to take antidiabetic and anticholesterol drugs than those in the WBD group
- There did not appear to be a trend for increasing volume of overweight/obese patients over the years

RESULTS




Figure 2. Discharge mRS of patients receiving weight-based dose vs max dose tPA




Figure 3. 90-d mRS of patients receiving weight-based dose vs max dose tPA

CONCLUSIONS

- **Overweight/obese patients who received MD IV tPA were more likely to have sICH and a worse discharge functional outcome vs. those in the WBD group**
- No difference in 90-d outcomes, likely resulting from a 28% lost to follow-up rate
- **Next step:** perform regression analysis to determine if the dosing itself explains the differences in sICH and outcomes
- The small n in the MD group did not allow for subgroup analyses; more cases will be added