

Baptist Health South Florida

## Scholarly Commons @ Baptist Health South Florida

---

All Publications

---

7-13-2019

### Implementation of an Inpatient Diabetes Team: Impact on Hypoglycemia Recurrence

Martha Bertin

*Homestead Hospital*, MarthaDa@baptisthealth.net

Elva Diaz

*Homestead Hospital*, ElvaD@baptisthealth.net

Follow this and additional works at: <https://scholarlycommons.baptisthealth.net/se-all-publications>



Part of the [Endocrine System Diseases Commons](#), [Nursing Commons](#), and the [Nutritional and Metabolic Diseases Commons](#)

---

#### Citation

Bertin, Martha and Diaz, Elva, "Implementation of an Inpatient Diabetes Team: Impact on Hypoglycemia Recurrence" (2019). *All Publications*. 3136.

<https://scholarlycommons.baptisthealth.net/se-all-publications/3136>

This Conference Poster -- Open Access is brought to you for free and open access by Scholarly Commons @ Baptist Health South Florida. It has been accepted for inclusion in All Publications by an authorized administrator of Scholarly Commons @ Baptist Health South Florida. For more information, please contact [Carrief@baptisthealth.net](mailto:Carrief@baptisthealth.net).

## Implementation of an Inpatient Diabetes Team: Impact on Hypoglycemia Recurrence

Maria M. Ojeda, DNP/PhD, MPH, BA, APRN, NP-C, BC-ADM<sup>1</sup>, Martha L. Bertin, DNP, APRN, FNP-BC<sup>2</sup>, Elva Diaz, BSN, RN, CDE<sup>2</sup>

<sup>1</sup>Walden University; <sup>2</sup>Homestead Hospital

### Introduction / Background

- Hypoglycemia among inpatients with diabetes (DM) can lead to complications, transfers to higher levels of care, increased length of stay, and mortality (Draznin et al., 2014; Tisminetsky et al., 2014).
- A substantial portion of hypoglycemic events are not followed by proper adjustments to medications or carbohydrate intake, leaving inpatients who experience hypoglycemia at high risk for recurrence (Cook et al., 2007; Maynard, Huynh, & Renvall, 2008). Some hospitals have instituted specialty teams to help manage patients with DM.

### Purpose

To evaluate the impact of a quality improvement program consisting of a DM-specialty team (nurse practitioner and RN-Certified Diabetes Educator) on rates of hypoglycemia recurrence during hospitalization.

### Aim


To describe and compare rates of hypoglycemia recurrence pre- and post-implementation of the DM-specialty team.

### References

- Cook, C. B., Castro, J. C., Schmidt, R. E., Gauthier, S. M., Whitaker, M. D., Roust, L. R., ... Zimmerman, R. S. (2007). Diabetes care in hospitalized noncritically ill patients: More evidence for clinical inertia and negative therapeutic momentum. *Journal of Hospital Medicine*, 2(4), 203-211. doi:10.1002/jhm.188
- Donabedian, A. (1988). The quality of care: how can it be assessed? *JAMA*, 260(12), 1743-1748.
- Draznin, B., Wang, Y., Seggelke, S., Hawkins, R. M., Gibbs, J., Bridenstine, M., ... Wang, C. L. (2014). Glycemic control and outcomes of hospitalization in non-critically ill patients with Type 2 diabetes admitted with cardiac problems or infections. *Endocrine Practice*, 20(12), 1303-1308. doi:10.4158/EP14174.DR
- Maynard, G., Huynh, M., & Renvall, M. (2008). Iatrogenic inpatient hypoglycemia: risk factors, treatment, and prevention. *Diabetes Spectrum*, 21(4), 241-247.
- Tisminetsky, M., Joffe, S., McManus, D. D., Darling, C., Gore, J. M., Yarzelski, J., ... Goldberg, R. J. (2014). Decade-long trends in the characteristics, management and hospital outcomes of diabetic patients with ST-segment elevation myocardial infarction. *Diabetes and Vascular Disease Research*, 11(3), 182-189. doi:10.1177/1479164114524235

### Theoretical Model

- Donabedian's (1988) *Structure, Process and Outcome* model for health care quality provided the framework for this study.
- The DM-specialty team was conceptualized as a structural element supporting the performance of care delivery processes linked to improved patient outcomes.




### Methods

- Design: Descriptive comparative.
- Sampling: Purposive sample of inpatients with DM who experienced at least 1-episode of hypoglycemia (blood glucose <70 mg/dl).
- Data Collection: Retrospective. Medical records of patients hospitalized during a 1-month period in 2015 prior to the implementation of the DM-specialty team, and those of patients seen by the DM-specialty team during a 1-month period in 2017, were extracted.
- Analysis: Data from 2015 was compared to 2017 on age and risk for mortality using a validated comorbidity index (Charlson Comorbidity Index). The rate of hypoglycemia recurrence (recorded episodes >15 minutes apart) was calculated.

### Results

- In 2015, 89 patients experienced hypoglycemia, in 2017 there were 71.
- The average age in 2015 was 65.5 years ( $sd=13.9$ ), in 2017 it was 63.4 years ( $sd=15.3$ ). The groups did not differ significantly on age ( $t(142)=.862$ ,  $p=.390$ ).
- Patients in 2017 had significantly higher comorbidity index scores ( $M=7.5$ ) compared to 2015 ( $M=5.6$ ), ( $t(120.217)=-2.820$ ,  $p=.006$ ).

Comparison of Hypoglycemia Rates



### Conclusions

- Patients seen by the DM-specialty team had substantially lower rates of hypoglycemia recurrence despite a significant increase in the risk for mortality associated with comorbidities.
- Quality and safety in the care of patients with DM may be improved through the addition of such teams within hospital settings.