

Baptist Health South Florida

Scholarly Commons @ Baptist Health South Florida

All Publications

10-25-2019

Preoperative factors related to delayed discharge after total knee arthroplasty removal from inpatient-only list

Juan Carlos Suarez

Baptist Health Medical Group; Miami Orthopedics & Sports Medicine Institute, juansu@baptisthealth.net

William Arguelles

Baptist Health South Florida, williamar@baptisthealth.net

Priscilla Rivera

Baptist Health South Florida, PriscillRi@baptisthealth.net

Yvette Hernandez

Baptist Health South Florida, yvettesal@baptisthealth.net

Don Parris

Baptist Health South Florida, donp@baptisthealth.net

See next page for additional authors

Follow this and additional works at: <https://scholarlycommons.baptisthealth.net/se-all-publications>

Citation

Suarez, Juan Carlos; Arguelles, William; Rivera, Priscilla; Hernandez, Yvette; Parris, Don; and Paraliticc, Giovanni, "Preoperative factors related to delayed discharge after total knee arthroplasty removal from inpatient-only list" (2019). *All Publications*. 3297.

<https://scholarlycommons.baptisthealth.net/se-all-publications/3297>

This Conference Poster -- Open Access is brought to you for free and open access by Scholarly Commons @ Baptist Health South Florida. It has been accepted for inclusion in All Publications by an authorized administrator of Scholarly Commons @ Baptist Health South Florida. For more information, please contact Carrief@baptisthealth.net.

Authors

Juan Carlos Suarez, William Arguelles, Priscilla Rivera, Yvette Hernandez, Don Parris, and Giovanni Paralicci

Preoperative Factors Related to Delayed Discharge after Total Knee Arthroplasty Removal from Inpatient-Only List

Juan C. Suarez, MD,¹ William Arguelles, PhD, MSPH,² Priscilla Rivera, MPH,² Yvette Hernandez,³ & Don Parris, PhD,²

¹ Miami Orthopedics & Sports Medicine Institute, Baptist Health South Florida; ² Center for Advanced Analytics, Baptist Health South Florida; ³ Center for Research, Baptist Health South Florida

Background

- The Centers for Medicare & Medicaid Services removed total knee arthroplasty (TKA) from the inpatient-only (IPO) list effective January of 2018
- Consequently, all TKAs were scheduled as outpatient (23-hour observation) without workflow modifications
- While previous studies have identified specific factors associated with successful outpatient TKAs, none have been described under these mandatory, non-selective circumstances¹⁻⁵
- Specific aim: to examine the preoperative factors associated with delayed discharge following this policy change among a non-selected cohort, as published data in this area are lacking

Methods

- TKA patients admitted to any Baptist Health South Florida facility between January 01, 2018 and August 01, 2018 were retrospectively analyzed
- Patients were stratified into two groups based on hospital night stay:
 - ≤ 1 night (non-delayed discharge)
 - > 1 night (delayed discharge)
- The association between the following preoperative factors and delayed discharge were examined using 2-tailed *t*-tests for continuous data, and Pearson's chi-square or Fischer exact tests for categorical data
- Age; gender; race/ethnicity; insurance type; preoperative education class attendance; anesthesia type; smoking; body mass index (BMI); diabetes; hypertension; cardiovascular diseases; chronic obstructive pulmonary disease (COPD); and kidney disease
- All analyses were conducted using SPSS (IBM, version 23.0)

Results

- 447 TKA patients were identified during the study period
 - 279 (62.4%) had a non-delayed discharge
 - 168 (37.6%) had a delayed discharge
- The following table presents the association between preoperative factors and delayed discharge among TKA patients originally scheduled as outpatient following TKA removal from IPO list

Variable	Nights ≤ 1 N = 279	Nights > 1 N = 168	P value
Age (years)	69.9 (8.6)	71.5 (8.0)	0.033
Gender			
Male	124 (44.4%)	39 (23.2%)	<0.001
Female	155 (55.6%)	129 (76.8%)	
Race/ethnicity			
Non-Hispanic White	107 (38.4%)	53 (31.5%)	0.348
Hispanic White	157 (56.3%)	105 (62.5%)	
Other or unknown	15 (5.4%)	10 (6.0%)	
Insurance type			0.185
Medicare HMO	97 (34.8%)	75 (44.6%)	
Medicare HCFA	107 (38.4%)	62 (36.9%)	
HMO	41 (14.7%)	15 (8.9%)	
PPO	20 (7.2%)	9 (5.4%)	
Other	14 (5.0%)	7 (4.2%)	
Preoperative education			0.782
Yes	61 (87.1%)	36 (83.7%)	
No	9 (12.9%)	7 (16.3%)	
Anesthesia type			0.111
General	60 (21.5%)	46 (27.4%)	
Spinal	79 (28.3%)	55 (32.7%)	
Spinal with MAC	135 (48.4%)	62 (36.9%)	
Other	5 (1.8%)	5 (3.0%)	
Smoking	29 (10.4%)	23 (13.7%)	0.291
BMI (kg/m ²)	30.8	31.2	0.464
Diabetes	33 (11.8%)	27 (16.1%)	0.252
Hypertension	109 (39.1%)	82 (48.8%)	0.049
Cardiac disease	4 (1.4%)	3 (1.8%)	1.000
CHF	1 (0.4%)	1 (0.6%)	1.000
Stroke	2 (0.7%)	1 (0.6%)	1.000
COPD	0 (0%)	4 (2.4%)	0.020
Kidney disease (creatinine > 1.5)	2 (0.7%)	0 (0%)	

Main Findings:

- Delayed discharge was related to:
 - Older age
 - Female sex
 - Hypertension
 - COPD
- Post-hoc analyses revealed that patients with delayed discharge were more likely to be discharged to an institution (26.8% vs. 5.8%, *p* < 0.001) than home (73.2% vs. 94.2%, *p* < 0.001), suggesting social support may also play an important role

Conclusions

- Almost 40% of our population had a ≥ 2 night hospital stay despite being scheduled as outpatient
- Preoperative factors associated with delayed discharge included older age, female sex, hypertension, and COPD
- With the numbers of available for study, delayed discharge was not related to race/ethnicity, insurance type, preoperative education, anesthesia type, smoking, body mass index, diabetes, cardiovascular disease, or kidney disease
- The influence of social support warrants further investigation since discharge to an institution was prevalent in the delayed group
- Limitations:**
 - Small sample size, with limited prevalence of certain conditions
 - Missing data on preoperative education class attendance

References

- Arshi, A., Leong, N. L., D'Oro, A., Wang, C., Buser, Z., et al. (2017). Outpatient total knee arthroplasty is associated with higher risk of preoperative complications. *The Journal of Bone and Joint Surgery*, 99, 1978-1986.
- Courtney, P. M., Friomson, M. I., Michael Meneghini, R. M., Lee, G., Della Valle, C. J., (2018). Can total knee arthroplasty be performed safely as an outpatient in the medicare population? *The Journal of Arthroplasty*, 1-4.
- Gauthier-Kwan, O. Y., Dobransky, J. S., Dervin, G.F., (2018). Quality of recovery, postdischarge hospital utilization, and 2-year functional outcomes after an outpatient total knee arthroplasty program. *The Journal of Arthroplasty*, 1-6.
- Hoffmann, J. D., Kusnezov, N. A., Dunn, J. C. Zarkadis, N. J., Goodman, G. P., et al. (2018). The shift to same-day outpatient joint arthroplasty: a systemic review. *The Journal of Arthroplasty*, 33, 1265-1274.
- Lovett-Carter, D., Sayeed, Z., Abaab, L., Pallekonda, V., Mihalko, W., et al. (2018). Impact of outpatient total joint replacement on postoperative replacement in postoperative outcomes. *Orthop Clin Am*, 49, 35-44.

Acknowledgements

- We'd like to thank Beth Willmitch, Zenia Reyes-Bardelas, Alicia d'Empaire, and their Information Technology teams for their data support

