

Baptist Health South Florida

Scholarly Commons @ Baptist Health South Florida

All Publications

10-25-2019

A Real World Experience of the FAST-ED Based Pre-Hospital Stroke Triage System to Detect Large Vessel Occlusions

Ned Shashoua

Miami Neuroscience Institute, neddavid12@gmail.com

Amy Starosciak

Miami Cancer Institute, amyst@baptisthealth.net

Jayme Strauss

Miami Neuroscience Institute, jaymes@baptisthealth.net

Felipe De Los Rios La Rosa

Miami Neuroscience Institute, felipedl@baptisthealth.net

Follow this and additional works at: <https://scholarlycommons.baptisthealth.net/se-all-publications>

Citation

Shashoua, Ned; Starosciak, Amy; Strauss, Jayme; and De Los Rios La Rosa, Felipe, "A Real World Experience of the FAST-ED Based Pre-Hospital Stroke Triage System to Detect Large Vessel Occlusions" (2019). *All Publications*. 3314.

<https://scholarlycommons.baptisthealth.net/se-all-publications/3314>

This Conference Poster -- Open Access is brought to you for free and open access by Scholarly Commons @ Baptist Health South Florida. It has been accepted for inclusion in All Publications by an authorized administrator of Scholarly Commons @ Baptist Health South Florida. For more information, please contact Carrief@baptisthealth.net.

INTRODUCTION

In March 2017, the Fire Officers Association of Miami-Dade (FOAMD) employed the Field Assessment Stroke Triage for Emergency Destination (FAST-ED) scale to better identify large vessel occlusion (LVO) strokes. Cases with a score ≥ 4 bypass other centers for a comprehensive center, and those ≥ 6 also have the interventional team activated from the field. The FAST-ED scale was developed using a retrospective set of ischemic strokes and TIAs. There have been no prospective, real world examples on FAST-ED performance when used in the field by emergency personnel.

Item	FAST-ED	NIHSS
Facial Palsy		
Normal or minor paralysis	0	0-1
Partial or complete paralysis	1	2-3
Speech Changes		
Absent	0	0
Mild to moderate	1	1
Severe, global aphasia, mute	2	2-3
Arm Weakness		
No drift	0	0
Limited drift, some resistance against gravity	1	1-2
No effort against gravity, absence of movement	2	3-4
Denial/Neglect		
Absent	0	0
Extinction to bilateral simultaneous stimulation in one sensory modality	1	1
Unable to recognize own hand, exclusively orients to only one side of the body	2	2
Eye Deviation		
Absent	0	0
Partial	1	1
Forced deviation	2	2

Table 1. FAST-ED Scale and Correspondence to NIHSS

METHODS

Stroke alert cases brought to our center by FOAMD during March 2017-August 2018 were analyzed. We used the FAST-ED score from the EMS run sheet and examined the incidence of LVOs and treatment rates for applicable cases. Cases without FAST-ED scores were excluded. An LVO was present if there was evidence of an occlusion of the internal carotid artery, middle cerebral artery (M1 or M2 segments), or basilar artery on initial imaging.




Figure 1. Number of Cases Reviewed

RESULTS




Figure 2. Stroke Types for FAST-ED ≥ 4 (A) and ≥ 6 (B)

- FAST-ED score ≥ 4 (Fig. 2A)
 - 60% were ischemic, 31% had an LVO
 - 23% were treated with mechanical reperfusion
- FAST-ED score ≥ 6 (Fig. 2B)
 - 66% were ischemic, 38% had an LVO
 - 31% were treated with mechanical reperfusion




Figure 3. Percent of LVOs **With** and **Without** MR Interventions of All Stroke Alerts per FAST-ED Score

CONCLUSIONS

- FAST-ED scores completed in the field yielded much lower LVO detection rates compared to the original report
- Likely due to inclusion of hemorrhages and stroke mimics from the field
- Only 1/3 of field activations led to an endovascular procedure
- Improved tools are required to identify stroke patients in the field with possible LVOs to minimize unnecessary interventionist field activations and provide appropriate destination triage

DISCLOSURES

None.

REFERENCES

1. Lima FO, Silva GS, Furie KL, et al. Stroke, 2016.