

Baptist Health South Florida

Scholarly Commons @ Baptist Health South Florida

All Publications

4-1-2021

Delayed Avascular Necrosis of the Talus Following Arthroscopic Os Trigonum Excision

Cary Chapman

Baptist Health Medical Group; Miami Orthopedics & Sports Medicine Institute, carych@baptisthealth.net

Follow this and additional works at: <https://scholarlycommons.baptisthealth.net/se-all-publications>

Citation

Foot & Ankle Orthopaedics (2021) 6(2):1-4

This Article -- Open Access is brought to you for free and open access by Scholarly Commons @ Baptist Health South Florida. It has been accepted for inclusion in All Publications by an authorized administrator of Scholarly Commons @ Baptist Health South Florida. For more information, please contact Carrief@baptisthealth.net.

Delayed Avascular Necrosis of the Talus Following Arthroscopic Os Trigonum Excision

Foot & Ankle Orthopaedics
2021, Vol. 6(2) 1-4
© The Author(s) 2021
DOI: 10.1177/24730114211015214
journals.sagepub.com/home/fao

Kshitij Manchanda, MD¹ , Brent J. Larson, DPM²,
and Cary B. Chapman, MD¹

Keywords: arthroscopy, avascular necrosis, osteonecrosis, Os trigonum, complications

Introduction

Posterior ankle arthroscopy is a commonly used modality for the treatment of os trigonum, prominent stieda process, posterior tibiotalar ligament impingement, removal of loose bodies, and tenolysis of the flexor hallucis longus.¹⁸ Complications following posterior ankle arthroscopy are rare but include damage to the posterior tibial artery and nerve when using the posterior medial portal.¹⁷

We present a case of iatrogenic avascular necrosis of the talus after arthroscopic excision of an os trigonum using a 2-incision posterior-medial and posterior-lateral approach. To our knowledge, this complication has not been reported after ankle arthroscopy.

Case

A 67-year-old healthy woman, with no significant medical or operative history, sustained a left inversion ankle sprain injury (no fractures on radiographic imaging) that was treated conservatively with rest, ice, compression, and elevation of the affected extremity. Four years following her injury, the patient continued to complain of posterior ankle pain that was aggravated with walking, navigating stairs, and wearing heels. Physical examination revealed tenderness and swelling of the posterior ankle joint. The patient was treated with a single cortisone injection, a nonsteroidal anti-inflammatory drug, physical therapy, and ankle supports without relief of symptoms. Radiographs and magnetic resonance imaging (MRI) revealed a large os trigonum involving the posterior talus with marrow edema within the posterior talus (Figure 1A-C). No additional corticosteroid injections were performed.

After 6 months of nonoperative treatment, the patient underwent a left posterior ankle arthroscopy with an excision of the os trigonum using the technique proposed by

van Dijk.¹⁶ The patient was in prone position and posterior arthroscopy was completed via posterior-medial and posterior-lateral portals with no distraction. The procedure was completed under tourniquet (total time of 40 minutes), with intra-articular fluid pressures maintained between 35 and 40 mm Hg. The flexor hallucis longus tendon was visualized and used as the medial boundary for dissection as the os trigonum was identified lateral to this. Soft tissue dissection medially was limited to avoid risking injury to neurovascular structures (Figure 2). The os was removed without complication and the patient's neurovascular status remained intact after surgery. The patient did well initially but developed pain approximately 6 months after surgery.

At 6 months following surgery, a plain-film radiograph demonstrated changes to the talus consistent with avascular necrosis (Figure 3). A follow-up MRI showed avascular necrosis (AVN) of the talar body (Figures 4 and 5). In the span of these 6 months, she did not sustain any new injuries or have any new changes in her medical or social history.

Discussion

Avascular necrosis of the talus has been reported following fracture-dislocation, exposure to pressurized atmospheric conditions, use of corticosteroids, Addison disease, Cushing syndrome, alcoholism, sickle cell, peripheral vascular

¹ Miami Orthopedics and Sports Medicine Institute, Coral Gables, FL, USA

² Richmond University Medical Center, Staten Island, NY, USA

Corresponding Author:

Cary B. Chapman, MD, Department of Orthopedic Foot and Ankle Surgery, 1228 South Pine Island Road, Ste. 310, Plantation, FL 33324, USA.

Email: CaryCh@baptisthealth.net






Figure 1. (A). Lateral plain film showing an os trigonum. (B) Coronal T1-weighted MRI showing no evidence of talar body injury. (C). Sagittal T2 MRI with effusion surrounding the os trigonum. MRI, magnetic resonance imaging.



Figure 2. Intraoperative arthroscopic view of os trigonum via posterior-lateral portal. Shaver is entering from posterior-medial portal and along the lateral aspect of the os and pointed toward the subtalar joint.



Figure 3. Lateral plain film of the foot after removal of the os trigonum.

disease, lupus, and dialysis for chronic renal failure.⁵ We report of a case of AVN of the talus that occurred after posterior ankle arthroscopy. Iatrogenic cases are rare and have been associated with clubfoot repair and triple arthrodesis. Two cases of avascular necrosis of the talus following posterior medial clubfoot approach are noted in the literature.² A case of AVN following a medial approach talonavicular arthrodesis has also been reported. These cases of iatrogenic AVN of the talus previously have been associated with damage to the artery of the tarsal canal.⁵

Posterior ankle arthroscopy is generally recognized as a safe procedure for removal of os trigonum, decompression of a prominent posterior talar process, tenolysis of flexor hallucis longus, removal of foreign bodies, and arthroscopy.¹⁸ The posterior medial portal is considered to be the highest risk but is found to be 16.5 mm from the posterior tibial artery.¹⁵

To our knowledge, no previous cases of AVN from ankle arthroscopy have been reported. However, there have been reported cases of AVN in other joints after arthroscopic procedures. Postarthroscopic osteonecrosis of the knee (or osteonecrosis in the postoperative knee) is a rare complication usually seen in elderly patients with recurrent or worsening pain after arthroscopic treatment of meniscal or chondroid lesions. The pathophysiology is still not fully understood but studies have suggested that it could be due to heat generated to the subchondral bone from radiofrequency probes or more likely from the increased biomechanical load causing insufficiency fractures within the femoral condyle after partial meniscectomy within the compartment.^{6,19} There have also been some case reports of AVN after arthroscopic anterior cruciate ligament reconstruction¹⁰ or patellar osteonecrosis after arthroscopy presumably due to excessive disruption of the anterior fat pad.³

Osteonecrosis of the femoral head with arthroscopy has been rarely seen and reported. The arthroscopy portals do not risk the key vascular structures, but possibly the combination of the traction and increased intracapsular pressure may increase the risk.^{13,14} Similarly, in the shoulder, few case



Figure 4. Sagittal T2-weighted magnetic resonance image of the talus, significant for avascular necrosis.




Figure 5. Coronal T1-weighted magnetic resonance image of the talus, significant for avascular necrosis.

series have reported postarthroscopic humeral head osteonecrosis after rotator cuff repairs. A potential cause of this has included the disruption of blood supply from the anchors.^{1,9}

AVN of the talus is primarily caused by disruption of the blood supply. In 1940, Phemister noted that the major blood supply to the talus was the dorsalis pedis artery and its branches.¹² The artery of the tarsal canal is another major blood supply to the body of the talus that arises from the posterior tibial artery. The peroneal artery supplies the lateral process of the talus. A significant portion of the blood supply lies in periosteal vessels around the talus.¹¹

Vasospasm has been shown to be an important mechanism in causing avascular necrosis. In a study on rabbits,

osteonecrosis was induced by injecting corticosteroids around the femoral head. In the treatment group, fasudil, an antispasmodic agent, was injected as well, and this significantly reduced the rate of avascular necrosis.⁷ A study on rats by Drescher et al⁴ also showed that vasoconstriction was a likely cause of femoral head avascular necrosis.⁷ Furthermore, patients with coagulopathy are predisposed to developing avascular necrosis.⁸ This all suggests that severance of arteries may not be necessary to cause this condition.

In this article, we have described a woman where the posterior medial and posterior lateral arthroscopic approaches were used for removal of an os trigonum. The MRI taken several months before the procedure failed to show any signs of avascular necrosis, and the MRI after the procedure demonstrates severe avascular necrosis of the talar body. The exact mechanism in this case is unclear as there was nothing to suggest arterial injury during or after surgery, nor was the case performed under tourniquet for an excessive period of time (40 minutes). Additionally, it seems she had no other extrinsic risk factors or causes that could have led to AVN. However, it is documented in animal models that vasoconstriction is a mechanism that can cause AVN. It would seem possible that in this case the arthroscopic procedure triggered a vasospastic event, causing avascular necrosis of the talus.

Ethics Approval

Ethical approval was not sought for the present study because this is a case report; however, we complied with HIPAA guidelines and maintained confidentiality and privacy while preparing the report.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

ORCID iD

Kshitij Manchanda, MD,  <https://orcid.org/0000-0002-1127-5312>

References

1. Beauthier V, Sanghavi S, Roulot E, et al. Humeral head osteonecrosis following arthroscopic rotator cuff repair. *Knee Surg Sports Traumatol Arthrosc.* 2010;18(10):1432-1434.
2. Blakeslee TJ, DeValentine SJ. Management of the resistant idiopathic clubfoot: the Kaiser experience from 1980–1990. *J Foot Ankle Surg.* 1995;34(2):167-176.
3. Carstensen SE, Domson GF. Patella osteonecrosis following knee arthroscopy. *Orthopedics.* 2019;42(6):e552-e554.
4. Drescher W, Lohse J, Varoga D, et al. Enhanced constriction of supplying arteries—a mechanism of femoral head Necrosis in Wistar rats? *Ann Anat.* 2010;192(1):58-61.

5. Hermus JPS. Osteonecrosis of the talus after talonavicular arthrodesis: a case report and review of the literature. *J Foot Ankle Surg.* 2011;50(3):343-346.
6. Higuchi H, Kobayashi Y, Kobayashi A, et al. Histologic analysis of postmeniscectomy osteonecrosis. *Am J Orthop (Belle Mead NJ).* 2013;42(5):220-222.
7. Ikemura S, Yamamoto T, Motomura G, et al. Preventive effects of the anti-vasospasm agent via the regulation of the Rho-kinase pathway on the development of steroid-induced osteonecrosis in rabbits. *Bone.* 2013;53(2):329-335.
8. Jones JP. (i) Coagulopathies in the pathogenesis of osteonecrosis. *Curr Orthop.* 1997;11(3):157-163.
9. Kim JK, Jeong HJ, Shin SJ, et al. Rapid progressive osteonecrosis of the humeral head after arthroscopic rotator cuff surgery. *Arthroscopy.* 2018;34(1):41-47.
10. Lansdown DA, Shaw J, Allen CR, et al. Osteonecrosis of the knee after anterior cruciate ligament reconstruction: a report of 5 cases. *Orthop J Sports Med.* 2015;3(3):2325967115576120.
11. Mulfinger GL, Trueta J. The blood supply of the talus. *J Bone Joint Surg Br.* 1970;52(1):160-167.
12. Phemister DB. Changes in bones and joints resulting from interruption of circulation. *Arch Surg.* 1940; 41(2): 436-472.
13. Scher DL, Belmont PJ Jr, Owens BD. Case report: osteonecrosis of the femoral head after hip arthroscopy. *Clin Orthop Relat Res.* 2010;468(11):3121-3125.
14. Sener N, Gogus A, Akman S, et al. Avascular necrosis of the femoral head after hip arthroscopy. *Hip Int.* 2011;21(5): 623-626.
15. Ürgüden M, Çevikol C, Kürşat Dabak T, et al. Effect of joint motion on safety of portals in posterior ankle arthroscopy. *Arthroscopy.* 2009;25(12):1442-1446.
16. van Dijk CN, Scholten PE, Krips R. A 2-portal endoscopic approach for diagnosis and treatment of posterior ankle pathology. *Arthroscopy.* 2000;16(8):871-876.
17. Voto SJ, Ewing JW, Fleissner PR Jr, et al. Ankle arthroscopy: neurovascular and arthroscopic anatomy of standard and trans-Achilles tendon portal placement. *Arthroscopy.* 1989;5(1): 41-46.
18. Willits K, Sonneveld H, Amendola A, et al. Outcome of posterior ankle arthroscopy for hindfoot impingement. *Arthroscopy.* 2008;24(2):196-202.
19. Zhuang Z, Chhantyal K, Shi Y, et al. Post-arthroscopic osteonecrosis of the knee: a case report and literature review. *Exp Ther Med.* 2020;20(4):3009-3016.